

Hidden Vector ‘Coordinates’ in Particle Physics

Gerald L. Fitzpatrick

PRI Research and Development Corp.

12517 131 Ct. NE

Kirkland, WA 98034

glfitzpatrick@yahoo.com

Abstract

Heretofore unrecognized (i.e., “hidden”) Lorentz-invariant vector *observables* in the fermion sector (i.e., flavor-defining fermion “coordinates”) are shown to indirectly explain most, if not all, of the so-called “accidental” (internal) symmetries associated with fundamental fermions (quarks and leptons), by explaining quark and lepton *flavors*, *flavor doublets* and *families*. Moreover, these new fermion coordinates lead to *quantitative* constraints on neutrino mixtures that are found to be in good agreement with current experimental observations. The new fermion coordinates arise in the context of a *quantized*, internal 2-space possessing a non-Euclidean metric. “Quantization” here means that all *scalars*, *vectors* and *matrices* relevant to a description of fundamental fermions in the 2-space, are both *discrete* and *finite* in number. In a certain limited sense, the new 2-space description of fundamental fermions is more fundamental, and more general, than that provided by the standard model of particle physics. In particular, it points to a “layer” of new physics—located *somewhere* between the unification scale, and the region of applicability of the standard model—that is responsible for the 2-space, its quantization, and its associated “selection rules.”

1.0 Introduction

Most physicists would probably agree that the standard model of particle physics [1–3], based on the gauge group $SU(3)_{\text{color}} \times SU(2)_L \times U(1)_Y$, is only a low-energy or “macroscopic” approximation to a deeper, more fundamental, *unified* theory, possibly involving either Planck level physics at an energy scale of 10^{19}GeV [4, 5], or a much lower TeV energy scale should the anticipated *extra* dimensions turn out to be large [6]. Consider for example, the case of the standard-model Lagrangian.

Given the number of flavors of quarks and leptons, and an appropriate (renormalizable) strong-electroweak Lagrangian, the so-called “accidental symmetries” of the Lagrangian [7] are known to “explain” both the existence, and separate (exact or approximate) conservation, of various (global) “charges” associated with quarks and leptons (e.g., *lepton number*, *baryon number*, *strangeness*, *charm*, *truth*, *beauty*, *electron-*, *muon-* and *tau-numbers*). However, *there is nothing in such Lagrangians, or their associated accidental symmetries, that would explain quarks and leptons, or tell us how many flavors of quarks and leptons to include in the Lagrangian*. Rather, it seems that the Lagrangian is *not* an explanation for these things, but instead is a *result* of physics at some deeper, more fundamental, level, possibly involving *unification*. However, if unification really does occur, we are immediately faced with the

important unanswered question of exactly how the standard model of particle physics, and its associated Lagrangian, arises or *emerges* from this deeper, more fundamental, “substratum.”

In this paper these questions are addressed in a new way by identifying what may turn out to be an intermediate “layer” of new physics located *somewhere* between the unification scale (wherever it happens to be), and the region of applicability of the standard model of particle physics. The strategy pursued here is similar to that of S. Goudsmit and G. Ulenbeck [8], who first realized that a description of the observed *fine structure* in atomic spectra was *not* possible unless electrons possessed, in addition to their mass and electric charge, a previously hidden *spin* “coordinate.” In particular, the principal purpose of this paper is to argue that the proposed intermediate “layer” of new physics is described by heretofore unrecognized (i.e., “hidden”) 2-vector observables in the fermion sector (i.e., flavor-defining fermion “coordinates” $\mathbf{U}$, $\mathbf{V}$ and $\mathbf{Q}$), which serve to explain, among many other things, the proliferation of matter fields (flavors) in comparison to the relative paucity of force-mediating particles. And, because we will also argue here that the new flavor-defining fermion coordinates (i.e., the 2-vectors $\mathbf{Q}$, $\mathbf{U}$, $\mathbf{V}$ where $\mathbf{Q} = \mathbf{U} + \mathbf{V}$) are *Lorentz-invariant*, we will be free to perform a variety of physically meaningful *algebraic*, *geometric*, and *topological* manipulations on any one (or more) of these coordinates—*without undue concern for how these manipulations “appear” to observers located in arbitrary (inertial) coordinate systems*.

From the foregoing, it is clear that once new coordinates of this kind are introduced in the fermion sector, an entirely new description of fundamental fermions (quarks and leptons) must result. Moreover, the proposed description appears to be both more fundamental, and more general (in certain limited respects), than the standard model description of fundamental fermions. For example, by employing these new fermion coordinates, the question of why fundamental fermion families are *replicated*, and the question of why there are just *three* families, can both be answered, whereas this is *not* possible in the standard model description [1–3]. Thus the new flavor-defining fermion coordinates serve both to *complement*, and to (modestly) *extend*, the standard model of particle physics.

This work is based, in part, on previous works by the author wherein different aspects of this approach to flavors, families, and the problem of family replication, were discussed [9–13]. These works provide a different emphasis, and in some cases, a more detailed description of many features of this general idea that are only touched upon in the present work, which essentially *reviews*, *clarifies*, and *consolidates* these earlier works.

2.0 The Electric Charge Coordinates $\mathbf{Q}$

It is well known that in quantum mechanics the wave function or state vector Ψ_n describing an individual (“isolated”) particle, provides all of the information that is available, in principle, about that particle. Here, the subscript on Ψ_n refers to various kinds of *simultaneously observable* quantum numbers associated with the particle. Hence, it is possible to effectively define such a quantum state by specifying the specific values of these simultaneous observables [14]. In a similar way, there are known to exist certain simultaneously observable flavor-defining quantum numbers (e.g., strangeness and baryon number) that

serve to define individual flavor or antiflavor eigenstates of quarks and leptons [15, 16]. It is proposed here that these flavor-defining quantum numbers are to be identified with certain global charge-conjugation-reversing (**C**-reversing) scalars, which are *derived*, in turn, from the flavor-defining fermion coordinates, **Q**, **U** and **V**.

If ever there was a signal for new coordinates in particle physics, surely it comes from the known fundamental-fermion sector. Unlike the known “force-mediating” fundamental-boson sector (i.e., the *photon*, *gluons*, the *graviton*, and *weak intermediate vector bosons*), matter-particles or fundamental fermions (quarks and leptons), exhibit a relative plethora of *flavors* falling into three families. Counting only color-flavor states, there are 48 known fundamental fermions and antifermions [17], compared with only 13 force-mediating bosons [18]. What could be more natural than to assume that this “embarrassment of riches” in the fermion sector, is due to new flavor-defining fermion coordinates, which are *not* carried by the force-mediating bosons [18, 19]?

In a recent paper [10], it was shown that the fermion-number operator **F**(op) itself may be the key to a preliminary understanding of how new flavor-defining fermion coordinates such as the electric-charge coordinate **Q** arise. In particular, let us review the salient points in [10] to see how these coordinates arise from an *analytic continuation* of the fermion-number operator **F**(op).

2.1 Analytic continuation of **F**(op) and a new internal non-Euclidean 2-space

We have found that an analytic-continuation of a Hermitian matrix **F**(op) representing the conventional fermion-number *operator*, from an *external* spacetime and Hilbert-space setting, to a new *internal* (real) non-Euclidean space—**F**(op) is continued to a real, generally non-diagonal matrix **F**(*v*) involving a single *real* parameter *v*—“automatically” leads to a new description of fundamental fermions (quarks and leptons) in which families are *replicated* and there are just *three* families. The fact that this happens, suggests that there is some deep connection between the result of the aforementioned analytic-continuation, namely **F**(*v*), and some more fundamental underlying physics (e.g., Planck-level physics) where flavor degrees-of-freedom, and family-replication, presumably originate.

2.1.1 The fermion-number operator **F**(op)

Consider the situation, presumably at some high energy, where we are dealing with “free” (isolated) leptons or “analytically free” quarks. Suppose we want to describe the scalar fermion-number carried by these particles. And, suppose further, that the energies involved are not so high that quantum field-theory (QFT) breaks down. Under these conditions the fermion-number can be represented by a $U(1)$ -type scalar “charge” [20, 21], namely, a charge associated with the (continuous) group of unitary matrices U of order 1 known as $U(1)$.

The fermion-number operator, which can be represented by a Hermitian matrix **F**(op), is said to generate these so-called “gauge” (or phase) transformations, which in turn act on

fermion and antifermion quantum states in Hilbert space. That is, given that α is a *real* phase one has

$$U = e^{i\alpha \mathbf{F}(\text{op})}, \quad (1)$$

and for infinitesimal transformations [i.e., $e^{i\delta\alpha \mathbf{F}(\text{op})} = 1 + i\delta\alpha \mathbf{F}(\text{op})$] acting on single-particle (free or “asymptotically free”) fermion and antifermion states $|p\rangle$ and $|\bar{p}\rangle$, respectively, one easily establishes that (the fermion-number “charges” are $f_m = -f_a = 1$ for matter and f_a for antimatter)

$$\begin{aligned} U|p\rangle &= e^{i\delta\alpha f_m}|p\rangle \\ U|\bar{p}\rangle &= e^{i\delta\alpha f_a}|\bar{p}\rangle, \end{aligned} \quad (2)$$

since, by definition, $\mathbf{F}(\text{op})$ obeys the eigenvalue equations

$$\begin{aligned} \mathbf{F}(\text{op})|p\rangle &= f_m|p\rangle \\ \mathbf{F}(\text{op})|\bar{p}\rangle &= f_a|\bar{p}\rangle. \end{aligned} \quad (3)$$

Finally, the assumption that the Hamiltonian H is invariant under U , namely

$$H = UHU^\dagger, \quad (4)$$

ensures that H and $\mathbf{F}(\text{op})$ commute

$$[\mathbf{F}(\text{op}), H] = 0, \quad (5)$$

as can be verified by differentiating $UHU^\dagger$ with respect to α . Hence, the total fermion-number (the number of *fermions* minus the number of *antifermions*) is a constant of the motion.

2.1.2 Matrix representation of the fermion-number operator $\mathbf{F}(\text{op})$

Because the matrix $\mathbf{F}(\text{op})$ involves just *two* kinds of quantum states (3), namely $|p\rangle$ and $|\bar{p}\rangle$, it can be expressed as a 2×2 diagonal Hermitian matrix (6), where one of the adjustable parameters (θ) is a *fixed* constant (up to 2π) and the other (ϕ) is freely *adjustable*. In particular,

$$\mathbf{F}(\text{op}) = \left(\begin{array}{cc} \cos \theta & \sin \theta e^{-i\phi} \\ \sin \theta e^{+i\phi} & -\cos \theta \end{array} \right) \Big|_{\cos \theta=1} = \sigma_z, \quad (6)$$

where

$$\sigma_z = \left(\begin{array}{cc} f_m & 0 \\ 0 & f_a \end{array} \right) \quad (7)$$

is one of the familiar Pauli matrices.

This form for $\mathbf{F}(\text{op})$ is consistent with (3) where the normalization and orthogonality conditions, namely $\langle p|p \rangle = \langle \bar{p}|\bar{p} \rangle = 1$ and $\langle p|\bar{p} \rangle = \langle \bar{p}|p \rangle = 0$, respectively, directly yield

$$\mathbf{F}(\text{op}) = \begin{pmatrix} \langle p|\mathbf{F}(\text{op})|p \rangle & \langle p|\mathbf{F}(\text{op})|\bar{p} \rangle \\ \langle \bar{p}|\mathbf{F}(\text{op})|p \rangle & \langle \bar{p}|\mathbf{F}(\text{op})|\bar{p} \rangle \end{pmatrix} = \sigma_z. \quad (8)$$

Note that owing to (7) and (8), $\cos \theta < 1$ in (6) is excluded. Here it should also be noted that $\text{tr} \mathbf{F}(\text{op}) = f_m + f_a = 0$, $\det \mathbf{F}(\text{op}) = f_m \cdot f_a = -1$, and $\mathbf{F}^2(\text{op}) = \mathbf{I}_2$ is the 2×2 identity matrix.

2.1.3 The Continuation From $\mathbf{F}(\text{op})$ to a ‘generalized fermion-number’ $\mathbf{F}(v)$

Now perform an analytic continuation on $\mathbf{F}(\text{op})$, namely $\mathbf{F}(\text{op}) \rightarrow \mathbf{F}(v)$, which maintains $\mathbf{F}(v)$ *real* and $\cos \theta \geq 1$. This can only be accomplished by continuing θ from a *real* to an *imaginary* number, and by maintaining $e^{-i\phi}$ *imaginary*. In particular, to maintain $\mathbf{F}(v)$ *real*, we must make the replacements $\theta \rightarrow iv$ and $e^{-i\phi} \rightarrow \mp i$, where v is a *real* number. Then

$$\mathbf{F}(v) = \left(\begin{array}{cc} \cos \theta & \sin \theta e^{-i\phi} \\ \sin \theta e^{+i\phi} & -\cos \theta \end{array} \right) \Big|_{\theta=iv, e^{-i\phi}=\mp i} \quad (9)$$

or

$$\mathbf{F}(v) = \left(\begin{array}{cc} \cosh v & \pm \sinh v \\ \mp \sinh v & -\cosh v \end{array} \right), \quad (10)$$

where, just as for $\mathbf{F}(\text{op})$, the eigenvalues of $\mathbf{F}(v)$ are f_m and f_a , and so we have $\text{tr} \mathbf{F}(v) = f_m + f_a = 0$, $\det \mathbf{F}(v) = f_m \cdot f_a = -1$, and $\mathbf{F}^2(v) = \mathbf{I}_2$.

In Ref. 9, pp. 50 and 54 it is argued that only the upper signs in (10) have physical significance and $v \geq 0$. And, just what is the physical significance of $\mathbf{F}(v)$?

Because the continuation “connects” $\mathbf{F}(\text{op})$ and $\mathbf{F}(v)$, it is natural to assume that *both* $\mathbf{F}(\text{op})$ and $\mathbf{F}(v)$ describe, or represent, aspects of the fermion number (i.e., the matter-antimatter “degree-of-freedom” or matter-antimatter “dichotomy”). However, unlike $\mathbf{F}(\text{op})$ or f , $\mathbf{F}(v)$ is a kind of “generalized fermion-number,” which will be shown to describe additional degrees-of-freedom such as the “up”-“down” and quark-lepton degrees-of-freedom. Moreover, it is abundantly clear from (10) that the generally non-Hermitian (when $v \neq 0$) matrix $\mathbf{F}(v)$ —unlike the Hermitian matrix $\mathbf{F}(\text{op})$ in (8), which acts on Hilbert space—does *not* act on a Hilbert space in an *external* spacetime setting [22].

2.1.4 A new internal non-Euclidean 2-space

Because of the matter-antimatter *dichotomy* exhibited by fundamental fermions, the fermion-number operator $\mathbf{F}(\text{op})$ is necessarily a 2 by 2 matrix, and so the continuation of $\mathbf{F}(\text{op})$, namely $\mathbf{F}(v)$, is also a 2 by 2 matrix. Therefore, *the matrix $\mathbf{F}(v)$ must “act” on some new internal 2-dimensional space*. What is the nature of this space? In particular, assuming that it is a metric space, what is the metric?

When the matrix $\mathbf{F}(v)$ acts on a real column-vector $\{a, b\}$, it leaves the quadratic form $a^2 - b^2$, invariant. Therefore, the 2-space metric is non-Euclidean or “Lorentzian”, and can be represented by the matrix [23]

$$\mathbf{g} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (11)$$

Given this metric, the *scalar product* of two real vectors (i.e., the “projection” of one vector upon another) assumes the form

$$(a, b)\{e, f\} = ae - bf. \quad (12)$$

Similarly, the *square* of a real vector (i.e., the “projection” of the vector upon itself) is given by

$$(a, b)\{a, b\} = a^2 - b^2. \quad (13)$$

In (12) and (13) above, $(\ , \)$ is a *row* vector while $\{ \ , \ }$ is a (conformable) *column* vector.

2.2 Quantization of the imaginary angle θ and the electric-charge coordinates $\mathbf{Q}$

It is very important to understand that the imaginary angle θ in (9) must be *quantized*. That is, when we do the continuation to an imaginary angle, the only allowed values seem to be

$$\theta = iv = i \ln M_c, \quad (14)$$

or

$$e^{-i\theta} = M_c, \quad (15)$$

where $M_c = 1$ or 3 only (see Refs. 24, 25, and Ref 9, p. 79).

This quantization of θ and/or v comes about in the following way. Because the eigenvalues of $\mathbf{F}(v)$ are $\mathbf{C}$ -reversing quantized *charges*, the components of both the diagonal and off-diagonal forms of the matrix $\mathbf{F}(v)$ are, necessarily, *quantized* charges. Since the components of $\mathbf{F}(v)$ involve hyperbolic functions of v , it follows that e^v , e^{-v} and v are, necessarily *quantized*. In particular, as we show elsewhere $fe^{-v} = f/M_c$ is a quantized charge identified with the *baryon* or *lepton* number (Ref. 9, pp. 52, 53), depending on the value of the quantum number M_c . And, from the spin-statistics theorem, we know that an *odd* number of fundamental fermions must bind together (“strongly”) to form a baryon or lepton. Hence, the quantum number M_c must be an odd integer $M_c = 1, 3, 5, \dots$ (see Refs. 24, 25, and Ref. 9, p. 79).




Figure 1. The *four* “electric charge” 2-vector fermion and antifermion “coordinates” $\mathbf{Q}_q, \mathbf{Q}_q^c, \mathbf{Q}_l, \mathbf{Q}_l^c$.

These identifications are remarkable for at least two reasons. First, we discover that the eigenvectors of $\mathbf{F}(v)$, and related functions evaluated at $M_c = 1$ and $M_c = 3$, yield the quantized *electric* charges of both leptons and quarks, and their associated quantized *lepton* or *baryon* numbers, respectively [26]. In particular, choose the upper signs in (10),

$$\mathbf{F}(v) = \begin{pmatrix} \cosh v & \sinh v \\ -\sinh v & -\cosh v \end{pmatrix}. \quad (16)$$

Then identify (see Ref. 9, pp. 52–55) the quark and lepton electric charges with the “up”-“down” components of the eigenvectors of the matrix $\mathbf{F}(v)$. That is, the quark charges are given by ($M_c = 3$)

$$q_1(f) = \frac{(M_c^2 - 1)}{2M_c(M_c - f)} = +\frac{2}{3} \text{ for } f = +1 \text{ and } +\frac{1}{3} \text{ for } f = -1, \quad (17)$$

$$q_2(f) = q_1(f) - 1, \quad (18)$$

where the baryon number for quarks is the non-Euclidean quadratic form $\mathbf{Q}_q^2 = B = q_1^2(f) - q_2^2(f) = \pm\frac{1}{3}$ for $f = \pm 1$, where the quark electric-charge “coordinate” (see Fig. 1) $\mathbf{Q}_q = \mathbf{Q}_q(f, M_c) = \{q_1(f), q_2(f)\}$ is an eigenvector of $\mathbf{F}(v)$. Similarly, the lepton electric charges

are given by ($M_c = 1$)

$$q'_1(f) = \frac{-(M_c^2 - 1)}{2M_c(M_c - f)} = -1 \text{ for } f = +1 \text{ and } 0 \text{ for } f = -1, \quad (19)$$

$$q'_2(f) = q'_1(f) + 1, \quad (20)$$

where the lepton number for leptons is the non-Euclidean quadratic form $\mathbf{Q}_l^2 = L = [q'_1(f)]^2 - [q'_2(f)]^2 = \pm 1$ for $f = \pm 1$, where the lepton electric-charge “coordinate” (see Fig. 1) $\mathbf{Q}_l = \mathbf{Q}_l(f, M_c) = \{q'_1(f), q'_2(f)\}$ is an eigenvector of $\mathbf{F}(v)$.

Second, it is abundantly clear that the quantum number M_c can be identified with the strong-color “multiplicity.” In particular, *leptons* being strong-color *singlets* ($M_c = 1$) do not bind strongly to form a composite particle, whereas *quarks* being strong-color *triplets* ($M_c = 3$), do bind strongly to form a composite baryon. That is, M_c is both a measure of the strong-color multiplicity and the *number* of quarks or leptons that bind (strongly) to form composite particles.

It is remarkable that the 2-space description of fundamental fermions has “automatically” yielded, not only the *electric*, *baric* and *leptic* charges of quarks and leptons, but also an unanticipated connection with *quantum chromodynamics* (QCD) and $SU(3)_{\text{color}}$. In particular, the 2-space and the associated quantized parameter $v = \ln M_c$, implies that something very similar to $SU(3)_{\text{color}}$ must be a symmetry of nature. *We take this circumstance to be a strong indication that the 2-space and/or whatever underlying physics is responsible for it, is more primitive (i.e., more fundamental) than the standard model of particle physics.*

2.2.1 Representing charge conjugation $\mathbf{C}$ in the 2-space

Given that there are numerous $\mathbf{C}$ -reversing scalars such as q_1, q_2, q'_1, q'_2, B and L , in the 2-space, there must exist a 2×2 matrix, call it $\mathbf{X}$, that serves to transform these *scalars*, various 2-*vector* fermion coordinates (e.g., $\mathbf{Q}, \mathbf{U}$ and $\mathbf{V}$), and various 2×2 *matrices* such as $\mathbf{F}(v)$, to their corresponding $\mathbf{C}$ -reversed (2-space) counterparts. For example, a matrix $\mathbf{X}$ should exist such that

$$\mathbf{X}\mathbf{Q} = \mathbf{Q}^c \quad (21)$$

and

$$\mathbf{X}\mathbf{Q}^c = \mathbf{Q}. \quad (22)$$

From (21) and (22) it follows that $\mathbf{X}$ must equal its multiplicative inverse ($\mathbf{X} = \mathbf{X}^{-1}$), and thus

$$\mathbf{X}^2 = \mathbf{I}_2, \quad (23)$$

where $\mathbf{I}_2$ is the 2×2 identity matrix.

Write $\mathbf{X}$ in the general form ($\mathbf{X}$ is real)

$$\mathbf{X} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad (24)$$

and consider the situation for leptons (see Eqs. 19 and 20), where one of the two charges (q'_2) associated with $\mathbf{Q}_l = \{q'_1(1) = -1 = q'_1, q'_2(1) = 0 = q'_2\}$ and $\mathbf{Q}_l^c = \{q'_1(-1) = 0 = q'_2, q'_2(-1) = +1 = -q'_1\}$ is *zero*, and the other charge (q'_1) is *nonzero*.

In this particular case, we have

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} q'_1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ -q'_1 \end{pmatrix}, \quad (25)$$

which means that

$$aq'_1 = 0 \quad (26)$$

and

$$cq'_1 = -q'_1. \quad (27)$$

And therefore, since $q'_1 \neq 0$, it must be true from (26) and (27) that $a = 0$ and $c = -1$.

Since $\mathbf{X}\mathbf{Q}^c = \mathbf{Q}$ it must also be true that

$$\begin{pmatrix} 0 & b \\ -1 & d \end{pmatrix} \begin{pmatrix} 0 \\ -q'_1 \end{pmatrix} = \begin{pmatrix} q'_1 \\ 0 \end{pmatrix}, \quad (28)$$

which means that

$$-bq'_1 = q'_1 \quad (29)$$

and

$$-dq'_1 = 0. \quad (30)$$

Finally, since $q'_1 \neq 0$ it must be true from (29) and (30) that $b = -1$ and $d = 0$.

Collecting the foregoing matrix elements, we have

$$\mathbf{X} = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \quad (31)$$

or

$$\mathbf{X} = -\boldsymbol{\sigma}_{\mathbf{X}}, \quad (32)$$

where $\boldsymbol{\sigma}_{\mathbf{X}}$ is one of the familiar Pauli matrices. In general, the matrix $\mathbf{X} = -\boldsymbol{\sigma}_{\mathbf{X}}$ should apply (in the 2-space) to 2-*scalars*, 2-*vectors*, 2×2 *matrices*, and to both *quarks* and *leptons*.

2.2.2 Transformations of charge-like scalars associated with $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$ under $\mathbf{X}$

As demonstrated in Section 2.2.1, the charge-like scalar *components* of 2-vector coordinates such as $\mathbf{Q}_q$ or $\mathbf{Q}_l$ change signs under $\mathbf{X}$. However, owing to the 2-space metric given by (11), scalar products of 2-vectors whose components transform like charges, will also be charge like.

For example, we have the *square* of the 2-vectors $\mathbf{Q}$ and $\mathbf{Q}^c$, namely,

$$\mathbf{Q}^2 = \mathbf{Q} \bullet \mathbf{Q} = q_1^2 - q_2^2 \quad (33)$$

and

$$(\mathbf{Q}^c)^2 = \mathbf{Q}^c \bullet \mathbf{Q}^c = q_2^2 - q_1^2. \quad (34)$$

Therefore,

$$\mathbf{Q}^2 = -(\mathbf{Q}^c)^2, \quad (35)$$

which means that $\mathbf{Q}^2$ and $(\mathbf{Q}^c)^2$ each transform like $\mathbf{C}$ -reversing 2-scalar *charges*.

As described previously, these particular charges can be identified with the *baryon-* or *lepton-number* carried by quarks or leptons, respectively (see Ref. 9, p. 72). We will see in a later section that when charge vectors such as $\mathbf{Q}$ (or $\mathbf{Q}^c$) are *resolved* in the 2-space into pairs of linearly independent vectors (e.g., $\mathbf{Q} = \mathbf{U} + \mathbf{V}$), not only are the *components* of $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$, $\mathbf{C}$ -reversing *charges*, but also by squaring $\mathbf{Q} = \mathbf{U} + \mathbf{V}$, namely,

$$\mathbf{Q}^2 = \mathbf{U}^2 + 2\mathbf{U} \bullet \mathbf{V} + \mathbf{V}^2, \quad (36)$$

$\mathbf{U}^2$, $2\mathbf{U} \bullet \mathbf{V}$ and $\mathbf{V}^2$ are, like $\mathbf{Q}^2$, $\mathbf{C}$ -reversing charges. The foregoing collection of 2-scalar charges will be used to define and describe *flavor eigenstates*, *flavor doublets*, and eventually *families* of fundamental fermions [27, 28].

2.2.3 Transformation of the metric and other matrices under $\mathbf{X}$

Any 2×2 matrix $\mathbf{M}$, appropriate to the 2-space description of fundamental fermions, should transform under $\mathbf{X} = -\boldsymbol{\sigma}_\mathbf{X}$ to its $\mathbf{C}$ -reversed counterpart $\mathbf{M}^c$ according to the *similarity transformation*

$$\mathbf{X} \mathbf{M} \mathbf{X}^{-1} = \mathbf{M}^c, \quad (37)$$

or because $\mathbf{X} = \mathbf{X}^{-1} = -\boldsymbol{\sigma}_\mathbf{X}$, equivalently as

$$(-\boldsymbol{\sigma}_\mathbf{X}) \mathbf{M} (-\boldsymbol{\sigma}_\mathbf{X}) = \mathbf{M}^c. \quad (38)$$

For example, the metric $\mathbf{g}$ (see Eq. 11) is found to be $\mathbf{C}$ -reversing since

$$(-\boldsymbol{\sigma}_\mathbf{X}) \mathbf{g} (-\boldsymbol{\sigma}_\mathbf{X}) = -\mathbf{g}. \quad (39)$$

Similarly, the matrix $\mathbf{F}(v)$ in (16) is **C**-reversing since $(-\sigma_{\mathbf{X}}) \mathbf{F}(v)(-\sigma_{\mathbf{X}}) = -\mathbf{F}(v)$.

A matrix that is **C**-invariant (e.g., the matrix $\mathbf{X}$) would, necessarily, have the form

$$\mathbf{N} = \begin{pmatrix} a & b \\ b & a \end{pmatrix}, \quad (40)$$

where it is clear that

$$(-\sigma_{\mathbf{X}})\mathbf{N}(-\sigma_{\mathbf{X}}) = \mathbf{N}. \quad (41)$$

3.0 What Dynamics Governs the Evolution of Flavor-Defining Fermion Coordinates $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$?

As a first approximation, it is safe to say that the *dynamics* of a system consisting of an “isolated” or individual elementary particle, is governed by the *evolution* of the system wave function or state vector. The state vector in turn evolves according to quantum mechanical equations of motion that can be said to arise from an appropriate strong-electroweak Lagrangian (when the standard model applies) or a more fundamental Lagrangian in case deeper physics applies [29]. The question that naturally arises in the present situation is how such a dynamical description is altered or constrained, if at all, by the new flavor-defining fermion coordinates $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$? Even a partial answer to this question may suggest ways that the new 2-space description of fundamental fermions can be experimentally tested. Let us begin the discussion by describing how quantum mechanical state vectors can be partially represented using the new 2-vector fermion coordinates $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$.

3.1 Representing flavor eigenstates and flavor doublets in the 2-space

The continuation from the 2-D Hilbert space to the 2-D non-Euclidean “charge” space (see Sec. 2.1.3) turns out to mean that individual flavors of fundamental fermions can be partially represented in an unconventional way by geometric objects (in the non-Euclidean 2-space) which *differ* from a quantum state, but *from which the quantum states can be inferred or effectively constructed*. In particular, in the non-Euclidean 2-space, an object we call a “vector triad” (i.e., the “triad” of flavor-defining fermion coordinates $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$) represents “up”-“down” type flavor *doublets* of fundamental fermions and antifermions—the “up”-“down” type flavor dichotomy. That is, the components of the vectors associated with a given vector-triad are *observable* Lorentz-invariant (Lorentz 4-scalar) “charges,” which can be used to define the *two* flavor-eigenstates in a flavor doublet [15, 27, 28, 16, 30].

3.1.1 Flavor eigenstates

As demonstrated in detail in Ref. 9, pp. 16–18, given the charge-like (**C**-reversing) *observables* associated with the description involving $\mathbf{F}(v)$, namely, the real **C**-reversing scalar-

components of various matrices and vectors defined on the internal non-Euclidean 2-space, it is possible to write down flavor eigenstates [16, 30].

In principle, what one does is to identify the mutually-commuting **C**-reversing “charges” (call them C_i) or charge-like quantum numbers associated with a particular flavor, and then write the corresponding simultaneous flavor-eigenstate as

$$|C_1, C_2, C_3, \dots, C_n\rangle. \quad (42)$$

Here $C_1, C_2, C_3, \dots, C_n$, are the “good” charge-like flavor-defining quantum numbers (charges) associated with a particular flavor. It happens that these *observable* real-numbers can be identified with quantum numbers such as *electric charge*, *strangeness*, *charm*, the third-component of (global) *isospin*, *truth and beauty* (see Ref. 9, p. 72). To discover what charges describe a particular flavor we must first identify the vector-triad associated with that flavor.

Now, each *vector-triad* represents a flavor *doublet*, not just an individual flavor. That is, vector-triads provide information on *two* quantum states (two simultaneous flavor-eigenstates) associated with flavor doublets. Therefore, vector-triads are associated with both individual flavors *and* flavor doublets. Here we simply summarize how it is that non-Euclidean vector-triads can be associated with both *individual* flavors and “up”-“down” type *flavor-doublets*.

3.1.2 Flavor doublets

Consider the eigenvector (call it **Q**) of $\mathbf{F}(v)$ for fundamental fermions (see Footnote 26, and Sec. 2.2). Since the space on which $\mathbf{F}(v)$ “acts” is two-dimensional, the *observable* vector **Q** can be “resolved” into two (no more or less) *observable*, linearly-independent vectors, call them **U** and **V**, as $\mathbf{Q} = \mathbf{U} + \mathbf{V}$ [31, 32]. Now, because these three vectors (**Q**, **U**, and **V**) are *simultaneous* observables, it makes sense to speak of this “triad” of vectors as being a well defined geometric object, namely, a “vector triad.”

Recognizing that the components of **Q**, **U** and **V** are **C**-reversing charge-like *observables* we can write these *observable* “charge” vectors as

$$\mathbf{Q} = \{q_1, q_2\} \quad (43)$$

$$\mathbf{U} = \{u_1, u_2\} \quad (44)$$

$$\mathbf{V} = \{v_1, v_2\}, \quad (45)$$

where q_1, q_2, u_1, u_2, v_1 and v_2 are the various *observable* “charges” (e.g., q_1 and q_2 were found to be *electric* charges). Given $\mathbf{Q} = \mathbf{U} + \mathbf{V}$, the non-Euclidean metric (11), and Eqs. (43) through (45), we find the associated *observable* quadratic “charges”

$$\mathbf{Q}^2 = \mathbf{U}^2 + 2\mathbf{U} \bullet \mathbf{V} + \mathbf{V}^2 \quad (46)$$

$$2\mathbf{U} \bullet \mathbf{V} = 2(u_1 v_1 - u_2 v_2) \quad (47)$$

$$\mathbf{U}^2 = u_1^2 - u_2^2 \quad (48)$$

$$\mathbf{V}^2 = v_1^2 - v_2^2. \quad (49)$$

Finally, using the foregoing flavor-defining charges, we can express the *two* quantum states (simultaneous flavor-eigenstates) associated with a *single* vector-triad in the form of “ket” vectors as follows (see Ref. 14, and Ref. 9, pp. 16–18)

$$\begin{aligned} &|q_1, u_1, v_1, \mathbf{Q}^2, \mathbf{U}^2, 2\mathbf{U} \bullet \mathbf{V}, \mathbf{V}^2\rangle, \\ &|q_2, u_2, v_2, \mathbf{Q}^2, \mathbf{U}^2, 2\mathbf{U} \bullet \mathbf{V}, \mathbf{V}^2\rangle. \end{aligned} \quad (50)$$

Here, the state $|q_1, u_1, v_1, \mathbf{Q}^2, \mathbf{U}^2, 2\mathbf{U} \bullet \mathbf{V}, \mathbf{V}^2\rangle$ represents the “up”-type flavor-eigenstate, and $|q_2, u_2, v_2, \mathbf{Q}^2, \mathbf{U}^2, 2\mathbf{U} \bullet \mathbf{V}, \mathbf{V}^2\rangle$ represents the corresponding “down”- type flavor-eigenstate in a flavor doublet of fundamental fermions [16, 30].

In this section we have demonstrated, among other things, that there are essentially two ways the new flavor-defining fermion coordinates $(\mathbf{Q}, \mathbf{U}, \mathbf{V})$ can be *combined*. In particular, when $(\mathbf{Q}, \mathbf{U}, \mathbf{V})$ apply to the *same* fundamental fermion, these vectors can either be *added*, as in $\mathbf{Q} = \mathbf{U} + \mathbf{V}$, or they can be *multiplied*, as in the non-Euclidean scalar products (i.e., “projections”) expressed by Equations 46–49. However, as described in the next section, *the linear superposition principle of quantum mechanics severely limits the ways $(\mathbf{Q}, \mathbf{U}, \mathbf{V})$ can be combined when two or more fundamental fermion flavors are involved*.

3.2 When the new internal coordinates *cannot* be superimposed

According to the linear superposition principle of quantum mechanics (QM), wave functions or state vectors for two or more particles, and certain of their associated simultaneously observable charge-like quantum numbers or “charges,” can be *superimposed*. And, because the overarching linear superposition principle of QM must be retained, it turns out that *like* flavor-defining fermion coordinates associated with (“carried by”) *two* or more fundamental fermions *cannot* be superimposed or *added* ($\mathbf{Q}$ with $\mathbf{Q}$, $\mathbf{U}$ with $\mathbf{U}$ and $\mathbf{V}$ with $\mathbf{V}$). For, if they could be superimposed, the linear superposition principle of QM could *not* be retained, nor could flavors be defined in a self-consistent way using the fermion coordinates $(\mathbf{Q}, \mathbf{U}, \mathbf{V})$.

In particular, one obvious reason that like coordinates $(\mathbf{Q}, \mathbf{U}$ or $\mathbf{V})$ for two or more fundamental fermions cannot be superimposed, is that such superpositions inevitably lead to nonsensical charge-like quantum numbers (“charges”), which could *not* be “carried” by a sensible state vector, or used to define the corresponding wave function or state vector for the collection of particles in question. Let us demonstrate the validity of this assertion by giving a few simple examples, which serve to demonstrate some of the rules governing the superposition (or lack thereof) of the 2-vector coordinates $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$.

3.2.1 Rules for combining $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$

Consider the case of three quarks bound together by strong forces to form a baryon with baryon number $B = 1$. Since the baryon number of each quark is given by $\mathbf{Q}^2 = 1/3$, the baryon number for three bound quarks is simply $3\mathbf{Q}^2 = B = 1$. Now this result makes it clear that we cannot simply add the three vectors $\mathbf{Q}$ together first, and then “square” them to get the baryon number for such a composite particle. In particular, it is clear that the baryon number of a composite consisting of three quarks is *not* given by $(3\mathbf{Q})^2 = B = 3$.

Now it might be supposed that this algorithmic failure can be attributed to the absence of an appropriate “normalization” factor when adding such vectors. For example, suppose that we were to make the rule that when N of these $\mathbf{Q}$ -vectors are added, we must divide the vector sum by $\sqrt{N}$. Then for the case of three quarks we would have a baryon number given by $(3\mathbf{Q}/\sqrt{3})^2 = B = 1$, which is certainly correct. However, the maximum, and minimum, electric charges of such a composite would then be given, respectively, by the scalar products $(3\mathbf{Q}/\sqrt{3})\{1, 0\} = +2\sqrt{3}/3$ and $(3\mathbf{Q}/\sqrt{3})\{0, -1\} = -\sqrt{3}/3$, where $\mathbf{Q} = (2/3, -1/3)$. Clearly, this is incorrect since three quarks with electric charges $+2/3$ obviously form a composite baryon with an electric charge of $+2$ not $+2\sqrt{3}/3$. Similarly, three quarks with electric charges $-1/3$ form a composite baryon with an electric charge of -1 not $-\sqrt{3}/3$. Moreover, if we choose not to employ the proposed normalization factor $\sqrt{N}$, it is true that the projections of the vector $3\mathbf{Q}$ upon the 2-space coordinate axes $\{1, 0\}$ and $\{0, -1\}$ do yield correct electric charges $+2$ and -1 as required [33]. However, in this case the baryon number of the composite is wrong, and there are no corresponding projection axes available for the *intermediate* electric charges between $+2$ and -1 that can be formed by combining three quarks in various well known ways. In particular, there is no way to obtain electric charges 0 and $+1$ by projecting the vector $3\mathbf{Q}$ upon the 2-space axes $\{1, 0\}$ or $\{0, -1\}$.

Considerations of the foregoing kind lead to the conclusion that *like* vector coordinates for two or more fundamental fermions *cannot* be added or superimposed ($\mathbf{Q}$ with $\mathbf{Q}$, $\mathbf{U}$ with $\mathbf{U}$, $\mathbf{V}$ with $\mathbf{V}$), the way quantum states or state vectors can be superimposed. Only the flavor-defining scalar *charges* associated with the flavor-defining coordinates $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$ for two or more fundamental fermions, can be combined or superimposed. That is, because flavor eigenstates are to be described in terms of various simultaneously-observable, global, flavor-defining (scalar) charge-like quantum numbers (associated with $\mathbf{Q}$, $\mathbf{U}$ or $\mathbf{V}$), it is simply *not* physically meaningful to superimpose like coordinates $\mathbf{Q}$, $\mathbf{U}$ or $\mathbf{V}$ for two or more particles [8].

The foregoing discussion clearly demonstrates that the dynamical significance of the new flavor-defining fermion coordinates $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$ is quite different than say, the intrinsic angular momentum or spin coordinates for two different fermions. Angular momentum vectors for two different particles can be combined, but the corresponding (like) $\mathbf{Q}$ -, $\mathbf{U}$ -, and $\mathbf{V}$ -vector coordinates cannot. Nevertheless, the triad of 2-vector coordinates ($\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$) used to represent a flavor eigenstate, is expected to change or *evolve* during certain kinds of *transitions* or *interactions*, simply because the wave function or state vector, which depends, in part, on flavor-defining global scalars associated with $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$, necessarily, *evolves*.

Clearly, the expectation that the vector triad *evolves* could have experimental consequences. As we will demonstrate in a later section, the evolution of vector triads, as *constrained* by new underlying physics, could lead to an explanation for, among other things, recent observations of solar neutrino mixing.

3.3 Quantization of the coordinates $\mathbf{U}$ and $\mathbf{V}$

Because the fermion-number operator $\mathbf{F}(\text{op})$ is a 2 by 2 matrix, its continuation (see Sec. 2.1.3), namely $\mathbf{F}(v)$, must likewise be a 2 by 2 matrix. Thus we have a natural explanation for why there must be *two* new linearly-independent flavor-defining fermion coordinates $\mathbf{U}$ and $\mathbf{V}$ associated with quarks and leptons, and their hypothetical superpartners [19]. Note that we have just *two* new (2-vector) coordinates $\mathbf{U}$ and $\mathbf{V}$ because the resultant 2-space can maintain at most two linearly independent 2-vector resolutions of $\mathbf{Q}$ at a time [31]. The question now arises as to how it is that these fermion coordinates can be *quantized*.

Since we do not have a first-principles understanding of the proposed quantization in terms of underlying dynamics, we will appeal to simple heuristic arguments that serve not only to place the question in perspective, but also yield *definite* rules for quantizing these coordinates.




Figure 2. This figure illustrates the *six* $\mathbf{U}$ -vector coordinates (solid arrows), and the *two* “electric charge” or $\mathbf{Q}$ -vector coordinates (dotted arrows) associated with *matter* particles (quarks q and leptons l), namely, $\mathbf{Q}_q$ and $\mathbf{Q}_l$. The horizontal “dotted” arrow is offset from the origin for purposes of illustration. It is assumed to pass through the origin.

3.3.1 Quantization of the $\mathbf{U}$ -vector

Write the 2-vectors $\mathbf{U}$ and $\mathbf{V}$ in terms of component charges, which we would like to quantize so as to quantize the coordinates $\mathbf{U}$ and $\mathbf{V}$. That is, write them as row $(\ , \)$ or column $\begin{pmatrix} \ \\ \ \end{pmatrix}$ vectors thus; $\mathbf{U} = (u_1, u_2)$ and $\mathbf{V} = (v_1, v_2)$ or $\mathbf{U} = \{u_1, u_2\}$ and $\mathbf{V} = \{v_1, v_2\}$, respectively. Now argue that *it should take only one charge type (i.e., u or v) to distinguish “up” and “down” type flavors* within any and all “up”-“down” type flavor doublets. If this is so, we can assume, without loss of generality, that this is the charge u (i.e., u_1 or u_2). Then, since the charge v does *not* distinguish “up” and “down” type flavors, it must be true that $v_1 = v_2$ for every flavor doublet of fundamental fermions. And, therefore, *given the metric in (11), the quantized charge $\mathbf{V}^2$ has but one value, namely, $\mathbf{V}^2 = (v_1, v_2) \bullet \{v_1, v_2\} = (v_1^2 - v_2^2) = 0$ for all fundamental fermions.* That is, all fundamental fermions are “neutral,” i.e., they are *singlets* with respect to the global charge $\mathbf{V}^2$.

Then, since $\mathbf{Q} = \mathbf{U} + \mathbf{V}$, we can write $\mathbf{Q}$ in column vector form in terms of $\mathbf{U}$ and $\mathbf{V}$ as $\mathbf{Q} = \{q_1, q_2\} = \{u_1, u_2\} + \{v_1, v_2\}$. Given that the difference between “up” and “down” electric charges for *all* fundamental fermions defines the fundamental (quantum) *unit* of electric charge (see Eqs. 18 and 20), namely, $q_1 - q_2 = \pm 1$, and given that $v_1 = v_2$ for all fundamental fermions, the vector $\mathbf{U}$ must have the column-vector form $\mathbf{U} = \pm\{a, a - 1\}$ for *all* fundamental fermions.

Now consider the nature of the charges u (i.e., u_1 and/or u_2), which are the components of $\mathbf{U} = \{u_1, u_2\}$. Because the charge u distinguishes “up” flavors from “down” flavors, and because it is a *charge*, it must be possible for this charge, like *all* known charges, to assume the value *zero*. Then, given that u -charge can be zero in certain cases, and given the form $\mathbf{U} = \pm\{a, a - 1\}$, we see immediately that the quantized values of a must include the values $a = 0$ or 1 . This means that there are at least four $\mathbf{U}$ -vectors, namely, the vectors $\mathbf{U} = \pm\{0, -1\}$ and $\mathbf{U} = \pm\{1, 0\}$. And, given the metric (11), this means that the quantized charge $\mathbf{U}^2$ must include the values $\mathbf{U}^2 = \pm 1$.

Finally, the quantized charge $\mathbf{U}^2$, like *all* known charges, should be able to assume the value *zero* for which $a = \frac{1}{2}$. Then the quantized values a are found to be limited to the *finite* sequence consisting of the *three* values $a = 0, \frac{1}{2}, 1$. Similarly, we find that the quantized values of the charge $\mathbf{U}^2$ are found to be limited to the *finite* sequence of *three* values $\mathbf{U}^2 = 0, \pm 1$, while the associated quantized $\mathbf{U}$ -vectors are limited to the *finite* sequence of *six* vectors (see Fig. 2), namely, $\mathbf{U} = \pm\{1/2, -1/2\}, \pm\{1, 0\}, \pm\{0, -1\}$.

3.3.2 Family replication and the number of families

Clearly, if we limit the discussion to matter particles—the same arguments apply to anti-matter particles and the vectors $\mathbf{Q}_q^c$ and $\mathbf{Q}_l^c$ —we note that because the vector $\mathbf{Q}_q$ applies to quarks and $\mathbf{Q}_l$ applies to leptons, some of the $\mathbf{U}$ - and $\mathbf{V}$ -vectors (and associated global charges) will apply to quarks, but *not* to leptons, and vice versa. So, it is appropriate to ask which of the six possible $\mathbf{U}$ -vectors go with the vector $\mathbf{Q}_q$, and which of the six possible $\mathbf{U}$ -vectors go with the vector $\mathbf{Q}_l$. Once this question is answered, one can easily determine which $\mathbf{V}$ -vectors go with the vector $\mathbf{Q}_q$, and which $\mathbf{V}$ -vectors go with the vector $\mathbf{Q}_l$. In

other words, we will be able to determine which $\mathbf{U}$ -vectors go with which $\mathbf{V}$ -vectors to form a vector triad $(\mathbf{Q}, \mathbf{U}, \mathbf{V})$ that serves to describe *flavor doublets* of quarks or leptons. This information will tell us, in turn, how many *generations* of quarks and leptons there are, and hence how many *families* of fundamental fermions there are.

By inspection of Figure 2 we see that *the requisite form for $\mathbf{V}$ -vectors, namely, $\mathbf{V} = \{v, v\}$ cannot be maintained if $\mathbf{U}$ -vectors in quadrant II (IV) are associated with the vector $\mathbf{Q}_q$ ($\mathbf{Q}_l$) in quadrant IV (II).* Therefore, we discover that the *single* vector $\mathbf{Q}_q$ ($\mathbf{Q}_l$) in quadrant IV (II), can only be associated with the *three* $\mathbf{U}$ -vectors located in the *same* quadrant. This simple result immediately tells us that, because there are only *six* $\mathbf{U}$ -vectors to choose from—three of which fall in quadrant IV, and three of which fall in quadrant II—there are only *three* generations of quark flavor doublets and *three* generations of lepton flavor doublets.

This is a remarkable result, for it means that a most unexpected explanation for why there are three families “automatically” emerges from the 2-space description. Of course, the underlying dynamics responsible for the 2-space, its quantization, and its associated selection rules would, if available, presumably provide a much more detailed explanation [29].

3.3.3 Quantization of the $\mathbf{V}$ -vector

Given the four $\mathbf{Q}$ -vectors (see Fig. 1), the six $\mathbf{U}$ -vectors from Section 3.3.1 and Figure 2, the associations between vectors such as $\mathbf{Q}_q$, $\mathbf{Q}_l$ and the six $\mathbf{U}$ -vectors (see Sec. 3.3.2), and the relation $\mathbf{Q} = \mathbf{U} + \mathbf{V}$, we have the following sequence of eleven quantized $\mathbf{V}$ -vectors, consistent with the requisite form $\mathbf{V} = \{v, v\}$, namely, $\mathbf{V} = \{0, 0\}$, $\pm\{1/6, 1/6\}$, $\pm\{-1/3, -1/3\}$, $\pm\{-1/2, -1/2\}$, $\pm\{2/3, 2/3\}$, $\pm\{-1, -1\}$ where the upper plus signs refer to matter, and the lower minus signs refer to antimatter. And, from the foregoing quantized $\mathbf{U}$ - and $\mathbf{V}$ -vectors we have the sequence of eleven quantized global charges $\mathbf{U} \bullet \mathbf{V} = 0, \pm 1/6, \pm 1/3, \pm 1/2, \pm 2/3, \pm 1$.

Here, the $\mathbf{V}$ -vectors $\pm\{1/6, 1/6\}$, $\pm\{-1/3, -1/3\}$ and $\pm\{2/3, 2/3\}$ apply to quarks (upper plus signs) and antiquarks (lower minus signs), while the $\mathbf{V}$ -vectors $\{0, 0\}$, $\pm\{-1/2, -1/2\}$ and $\pm\{-1, -1\}$ apply to leptons (upper plus signs) and antileptons (lower minus signs). Similarly, the charges $\mathbf{U} \bullet \mathbf{V} = \pm 1/6, \mp 1/3, \pm 2/3$ apply to quarks (upper signs) and antiquarks (lower signs), while the charges $\mathbf{U} \bullet \mathbf{V} = 0, \mp 1/2, \mp 1$ apply to leptons (upper signs) and antileptons (lower signs). Note that the scalar components of $\mathbf{V}$ -vectors are proportional to the associated charges $\mathbf{U} \bullet \mathbf{V}$.

3.3.4 A natural basis for family membership of flavor doublets

The discussion in Section 3.3.2 makes it clear that there are only *three* quark-lepton families. But, what determines which *quark* flavor doublet goes with which *lepton* flavor doublet to make a particular quark-lepton *family*? In other words, how does one decide which quark $\mathbf{U}$ -vector goes with which lepton $\mathbf{U}$ -vector?

The question of family membership of flavor doublets is *not* answered in the standard model of particle physics where these family assignments are made, more-or-less, on the basis of the mass hierarchy of the average family member. However, in the 2-space, there are

other, more “natural,” ways of assigning flavor doublets to families (see Ref. 9 for a detailed discussion). We note first the fact that two of the three *quark* flavor doublets associated with the three $\mathbf{U}$ -vectors $\mathbf{U} = \{1/2, -1/2\}$, $\{1, 0\}$ and $\{0, -1\}$, are “equidistant” from the “intermediate” $\mathbf{U}$ -vector $\{1, 0\}$, with respect to a matrix $\mathbf{R}$ given by

$$\mathbf{R} = \begin{pmatrix} 0 & -2 \\ -1 & -1 \end{pmatrix}. \quad (51)$$

That is, the matrix $\mathbf{R}$ is a Lorentz-invariant “measure” of the “distance” between certain $\mathbf{U}$ -vectors, and their associated flavor doublets, thus [34, 35]

$$\{1/2, -1/2\} \xrightarrow{\mathbf{R}} \{1, 0\} \xrightarrow{\mathbf{R}} \{0, -1\}. \quad (52)$$

Similarly, two of the three *lepton* flavor doublets associated with the three $\mathbf{U}$ -vectors $\{-1/2, 1/2\}$, $\{-1, 0\}$ and $\{0, 1\}$, are also “equidistant” from the “intermediate” $\mathbf{U}$ -vector $\{-1, 0\}$, with respect to the *same* matrix $\mathbf{R}$ since

$$\{-1/2, 1/2\} \xrightarrow{\mathbf{R}} \{-1, 0\} \xrightarrow{\mathbf{R}} \{0, 1\}. \quad (53)$$

In other words, the $\mathbf{U}$ -vector *pairs* $[\{1/2, -1/2\} \text{ and } \{-1/2, 1/2\}]$ or $[\{1, 0\} \text{ and } \{-1, 0\}]$ or $[\{0, -1\} \text{ and } \{0, 1\}]$ are at the *same* “distance” or “location” with respect to the single “distance” measure $\mathbf{R}$. That is, they are associated with, or are “in,” the *same* family. To state this differently, *the $\mathbf{U}$ -vector describing a quark flavor doublet in a given family is the additive inverse of the $\mathbf{U}$ -vector describing a lepton flavor doublet in the same family, and vice versa.*

Because of the foregoing relations and identifications, it would seem very natural to assign the $\mathbf{U}$ -vectors $\pm\{1/2, -1/2\}$ to the *first* family, $\pm\{1, 0\}$ to the *second*, or intermediate family, and $\pm\{0, -1\}$ to the *third* family, where the upper + signs refer to *quarks* and the lower – signs refer to *leptons*. While the assignment of the $\mathbf{U}$ -vectors $\pm\{1, 0\}$ to the *second*, or intermediate, family seems unambiguous, what prevents us from assigning the $\mathbf{U}$ -vectors $\pm\{1/2, -1/2\}$ to the *third* family, and the $\mathbf{U}$ -vectors $\pm\{0, -1\}$ to the *first* family?

After all, we could just as well have used the multiplicative *inverse* of the matrix $\mathbf{R}$, namely, $\mathbf{R}^{-1}$ where

$$\mathbf{R}^{-1} = \begin{pmatrix} \frac{1}{2} & -1 \\ -\frac{1}{2} & 0 \end{pmatrix}, \quad (54)$$

to be the Lorentz-invariant measure of “distance,” in which case we would have had the following relations and associations

$$\pm\{1/2, -1/2\} \xleftarrow{\mathbf{R}^{-1}} \pm\{1, 0\} \xleftarrow{\mathbf{R}^{-1}} \pm\{0, -1\}. \quad (55)$$

Clearly, from these relations and associations it would seem just as natural to assign the $\mathbf{U}$ -vectors $\pm\{1/2, -1/2\}$ to the *third* family, $\pm\{1, 0\}$ to the *second*, or intermediate family,

and $\pm\{0, -1\}$ to the *first* family. How then, are we to decide if the $\mathbf{U}$ -vectors $\pm\{1/2, -1/2\}$, are to be assigned to the *first* or the *third* family? The 2-space description of quarks and leptons provides a simple “answer” to this question as well.

It turns out (see Ref. 9, pp. 56–58) that the vector triads $(\mathbf{Q}, \mathbf{U}, \mathbf{V})$ associated with both quarks and leptons have the *highest* possible *degree* of symmetry with respect to the matrix transformation $\mathbf{F}(v)$, *only* when the associated $\mathbf{U}$ -vectors are $\pm\{1/2, -1/2\}$. And, because the “ground state” is typically the most probable state, with the highest degree of symmetry into which other less symmetrical states tend to *evolve* [36], it is natural to assume that the $\mathbf{U}$ -vectors $\pm\{1/2, -1/2\}$ are associated with the *first* or “ground state” family. Recall that the members of the first family have the lowest average mass of any family members, and so the first family naturally resembles a kind of “ground state.”

The foregoing associations and identifications, together with the specific family assignments of the *experimentally* observed quarks and leptons, are elaborated elsewhere (see Ref. 9). However, it should be clear—as we have repeatedly emphasized throughout this paper—that the foregoing heuristic 2-space relations and associations must be a consequence of some “deeper,” more fundamental, “layer” of physics that is responsible for the 2-space, its quantization, and its associated selection rules.

3.4 The nature of transitions of vector triads

Because strong and electromagnetic interactions have no effect on flavor indices, they can have no effect on the flavor-defining fermion coordinates. Hence, vector triads $(\mathbf{Q}, \mathbf{U}, \mathbf{V})$ will be unaffected by *strong* and *electromagnetic* interactions. However, in general, this is not the case for *weak* interactions.

Because the quantum state of an “isolated” fundamental fermion such as a quark (eventually) evolves via weak interactions, so must the associated vector triad, consisting of the three 2-vectors $(\mathbf{Q}, \mathbf{U}$ and $\mathbf{V})$ *evolve*, and vice versa. And, if we knew the dynamics [29] underlying the 2-space, presumably we would be in a position to calculate, from first principles, such things as Cabibbo-Kobayashi-Maskawa (CKM) matrix elements. However, as this dynamics is currently unknown, we will have to satisfy ourselves here with a less ambitious description based on mathematical properties of the 2-space, and certain experimental facts about transitions of fundamental fermions. Consider first the so-called “diagonal” weak transitions of quarks.

Diagonal weak transitions of quarks are “up”-“down” or “down”-“up” type transitions involving the two quark flavors (flavor eigenstates) “located” within the *same* flavor doublet. From experimental data on CKM matrix elements, these diagonal transitions are known to be far more probable than the so-called “off-diagonal” transitions, which are relatively suppressed. What do these experimental facts teach us about the *evolution* of vector triads?

In the case of diagonal transitions, it is clear that the vector triad associated with the *initial* quark state, and the vector triad associated with the *final* quark state, are identical, i.e., they are *congruent*. By comparison, the initial and final state triads associated with all “off-diagonal” weak transitions of quarks are *different*. These observations suggest that

the underlying dynamics tends to “prefer” transitions in which the associated vector triad is *unchanged*. Clearly, such transitions would be characterized by the “selection rules” (use $\mathbf{Q} = \mathbf{U} + \mathbf{V}$)

$$\Delta\mathbf{Q} = \Delta\mathbf{U} = \Delta\mathbf{V} = 0. \quad (56)$$

Moreover, since $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$ don’t change in such transitions, we also have the selection rules

$$\Delta\mathbf{Q}^2 = \Delta\mathbf{U}^2 = \Delta\mathbf{U} \bullet \mathbf{V} = 0, \quad (57)$$

which means that the baryon number $B = \mathbf{Q}^2$, the charge $\mathbf{U}^2$, and the charge $\mathbf{U} \bullet \mathbf{V}$, are separately *maintained* or *conserved* in such transitions [37].

While quark masses play a significant role in determining the magnitudes of CKM matrix elements, there appears to be a trend here, namely, this: “Easy” transitions (i.e., diagonal transitions *within* a quark generation) are those for which the associated vector triad does *not* change, and “hard” transitions (i.e., off-diagonal transitions *between* different quark generations) are those for which it *does* change.

To be more specific, while there are some 24 flavor and antiflavor eigenstates of quarks and leptons [17], there are just 12 *distinct* vector triads. That is, the vector triad associated with each flavor doublet is associated with each of *two* flavor or antiflavor eigenstates. This means that all diagonal transitions involve the *same* vector triad, and all off-diagonal transitions involve *different* (distinguishable) vector triads.

What is it that differs from vector triad to vector triad? Of course, charge-like quantum numbers such as

$$\mathbf{Q}^2, \mathbf{U}^2, \text{ and } \mathbf{U} \bullet \mathbf{V} \quad (58)$$

can sometimes differ. But there are other, more subtle, things that can differ as well. It happens that different vector triads, and certain of their associated 2-vector coordinates, are “isolated” from one another in a *topological* sense. Take for example the four 2-vectors $\mathbf{Q}_q$, $\mathbf{Q}_q^c$, $\mathbf{Q}_l$, and $\mathbf{Q}_l^c$ illustrated in Figure 1.

Because there are no *continuous* transformations (associated with the metric in Eq. 11) available to convert the 2-vectors $\mathbf{Q}_q$, $\mathbf{Q}_q^c$, $\mathbf{Q}_l$ and $\mathbf{Q}_l^c$ one into the other, these four $\mathbf{Q}$ -vectors are, necessarily, *topologically distinct*. It is certainly conceivable that *this fact could be an important underlying reason for the separate conservation of baryon number B and lepton number L* in most physical processes. In particular, because the $\mathbf{Q}$ -vectors are *topologically distinct*, underlying dynamics could conceivably dictate that there are topological energy “barriers” that *maintain* these topological distinctions, leading to the 2-vector selection rule [39]

$$\Delta\mathbf{Q} = 0, \quad (59)$$

and consequently to the selection rule (Note that $\mathbf{Q}^2 = B$ or L)

$$\Delta\mathbf{Q}^2 = 0, \quad (60)$$

which should then apply to *all* fundamental fermions [39].

Of course, the discussion in the previous paragraph involves only *one* of the three 2-vector coordinates (i.e., $\mathbf{Q}$) associated with a vector triad $(\mathbf{Q}, \mathbf{U}, \mathbf{V})$. There are other topological distinctions that can be drawn, which involve all *three* 2-vectors associated with a particular vector triad. For example, as shown elsewhere (see Ref. 9, pp. 56, 57, and Appendix A in Ref. 13), and as elaborated in the following section, there are also topological distinctions based on how an *entire* vector triad $(\mathbf{Q}, \mathbf{U}, \mathbf{V})$ *transforms* under the matrix $\mathbf{F}(v)$. It happens that some vector triads transform like Möbius strips, and some transform like cylinders under $\mathbf{F}(v)$. Clearly, if there were underlying topological energy “barriers” that act to inhibit changes in the topology of *entire* vector triads, this situation could have experimental consequences [38]. And, if experiments were to eventually confirm that such topological energy “barriers” exist, a new level of dynamics “below” or “beyond” the standard model of particle physics would seem to be required to explain them.

3.5 Does the dynamics underlying the 2-space impose topological constraints on 3-flavor neutrino mixtures?

As alluded to in the previous section, it is important to understand that the new 2-space description of fundamental fermions (quarks and leptons) provides a distinction between these particles that goes beyond differences that can be explained by mass differences alone. For example, in the standard model of particle physics the only difference between the u , c and t quarks is that they have different masses. Otherwise, these particles experience identical strong and electroweak interactions. Moreover, the separate conservation (in strong and electromagnetic interactions) of quantum numbers such as “charm” and “truth” can be attributed to certain unavoidable “accidental symmetries” associated with the (renormalizable) Lagrangian describing the interactions of these particles [7].

Taken at face value, these accidental symmetries would seem to imply that there are no internal “wheels and gears” that would distinguish a u quark from a c quark, for example. But, if the string theories are correct, these particles would be associated with different “handles” on the compactified space (see Ref. 4, Vol. 2, p. 408), and so would be different in this *additional* sense. Likewise, in the present non-Euclidean 2-space description, *topological* differences in addition to a variety of (global) 2-scalars, which are *indirectly* related to the accidental symmetries of the Lagrangian, serve to provide further distinctions between matter particles.

A possible experimental signal of such internal differences is to be found in the recent observations at the Super Kamiokande and SNO of (nearly) bi-maximal $\nu_\mu - \nu_\tau$ neutrino mixing [43]. Models which begin by positing a neutrino mass-matrix, and associated mixing-parameters, such as the three-generation model proposed by Georgi and Glashow [40], do an acceptable job of describing the observations. However, (nearly) bi-maximal $\nu_\mu - \nu_\tau$ mixing may have a deeper explanation in terms of internal topological differences (in the non-Euclidean 2-space) between ν_e , and ν_μ or ν_τ neutrinos.

With respect to the internal transformation $\mathbf{F}(v)$, the topology of the non-Euclidean

“vector triad” (see Ref. 9, p. 57, and the qualifying remarks in Appendix A of Ref. 13) associated with the ν_e (ν_μ or ν_τ), is found to be that of a cylinder (Möbius strip). And, *assuming that a change in topology (of vector triads) during neutrino mixing is suppressed by energy “barriers,” or other topological “barriers”* described by physics “beyond” the standard model, while neutrino mixing without topology-change is (relatively) enhanced [38, 41, 42], one can readily explain the experimental observation of (nearly) bi-maximal $\nu_\mu - \nu_\tau$ neutrino mixing—at least (nearly) bi-maximal $\nu_\mu - \nu_\tau$ mixing over long distances, where the proposed topological influences are expected to be *cumulative*. If this qualitative explanation (see detailed calculations below) is basically correct, then it follows that the neutrino mass-matrix, and associated mixing-parameters needed to explain (nearly) bi-maximal $\nu_\mu - \nu_\tau$ neutrino mixing, would be the *result* of new physics (i.e., dynamics) associated with these deeper (internal) topological differences between neutrinos, and not their *cause*.

3.5.1 Conventional description of 3-flavor neutrino mixing

Except where explicitly prevented by some “absolute” conservation law (e.g., the conservation of electric charge or spin angular momentum), quantum mechanics generally permits transitions between states having *different* topologies [41]. While a change in topology may be energetically (or otherwise) inhibited by underlying dynamics, unavoidable quantum fluctuations are expected to *catalyze* such processes. Hence, there is always the possibility of *mixing* between otherwise similar quantum states (e.g., ν_e , ν_μ and ν_τ) associated with vector triads having *distinct* topologies [42].

As described previously—with respect to the matrix transformation $\mathbf{F}(v)$ —the topology associated with both the ν_μ and ν_τ is found to be that of a Möbius strip (Ref. 47, p. 143). By contrast, the topology associated with the ν_e [with respect to $\mathbf{F}(v)$] is that of a cylinder. And because it is reasonable to *assume* that a change in topology during transitions tends to be *suppressed* [38], the foregoing topological distinctions between neutrinos may help explain recent observations of (nearly) bi-maximal $\nu_\mu - \nu_\tau$ mixing [43]. However, it is important to stress at the outset that there are good reasons for believing that similar topological distinctions (alone) among quarks, while also present, do *not* play an important role in d , s , b quark mixing [44].

At “birth” via weak decays, or upon detection via weak capture interactions, neutrinos have a *definite* flavor and associated topology [45]. However, between birth and detection they are in a mixed state having no definite flavor or associated topology. In this intermediate region the probability of flavor (or topology) maintenance and/or change *oscillates*. Only at great distances from the neutrino source do these oscillations finally “damp out.”

In the conventional description of three-flavor neutrino mixing *flavor eigenstates* are related to neutrino *mass eigenstates* (states of definite mass) via a unitary CKM-like “mixing” matrix $U_{\alpha i}$, as follows (Ref. 3, p. 365 and Ref. 46)

$$\nu_\alpha = \sum_{i=1}^3 U_{\alpha i} \nu_i. \quad (61)$$

Here, $\nu_i = \nu_1, \nu_2$ and ν_3 are mass eigenstates with mass eigenvalues m_1, m_2 and m_3 , respectively, while $\nu_\alpha = \nu_e, \nu_\mu$ and ν_τ are flavor eigenstates. Using a conventional parameterization for $U_{\alpha i}$, (61) becomes

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} c_1 & s_1 c_3 & s_1 s_3 \\ -s_1 c_2 & c_1 c_2 c_3 - s_2 s_3 e^{i\delta} & c_1 c_2 s_3 + s_2 c_3 e^{i\delta} \\ -s_1 s_2 & c_1 s_2 c_3 + c_2 s_3 e^{i\delta} & c_1 s_2 s_3 - c_2 c_3 e^{i\delta} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}, \quad (62)$$

where $c_i \equiv \cos \theta_i$ and $s_i \equiv \sin \theta_i$, and δ is associated with a Dirac-type CP-noninvariant phase factor $e^{i\delta}$.

Using (62) it can be shown that the probability of detecting a neutrino of flavor type β at a distance X from a source of neutrinos of flavor type α is given by [46]

$$P_{\nu_\alpha \rightarrow \nu_\beta} = \sum_{i=1}^3 |U_{\alpha i}|^2 |U_{\beta i}|^2 + \sum_{i \neq j}^3 U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j} \cos \left(\frac{2\pi X}{l_{ij}} \right). \quad (63)$$

Here, the so-called *oscillation lengths* l_{ij} are given by $l_{ij} = 2\pi/(E_i - E_j)$, where the total relativistic energy differences in a beam of neutrinos having *fixed* momentum p are $E_i - E_j = (m_i^2 - m_j^2)/2p$.

Examination of (63) shows that at the neutrino source ($X = 0$, and $t = 0$) the probability $P_{\nu_\alpha \rightarrow \nu_\beta}$ reduces, as expected, to the 3×3 identity matrix

$$P_{\nu_\alpha \rightarrow \nu_\beta} \Big|_{X=0} = I_3, \quad (64)$$

while at “intermediate” distances from the neutrino source ($X \approx l_{ij}$), the probability $P_{\nu_\alpha \rightarrow \nu_\beta}$ undergoes *oscillations*. Finally, at great distances from the neutrino source ($X \gg l_{ij}$), all time- or distance-dependent oscillations “damp out,” and we are left with a 3×3 matrix of time-average probabilities [46], namely,

$$\langle P_{\nu_\alpha \rightarrow \nu_\beta} \rangle = \sum_{i=1}^3 |U_{\alpha i}|^2 |U_{\beta i}|^2. \quad (65)$$

From (65) it is clear that this matrix (α = row index, β = column index), which describes long-distance neutrino mixtures, is *symmetric*. Keeping in mind this symmetry, and calling this matrix M , one has

$$M = \begin{pmatrix} \langle P_{\nu_e \rightarrow \nu_e} \rangle & \langle P_{\nu_e \rightarrow \nu_\mu} \rangle & \langle P_{\nu_e \rightarrow \nu_\tau} \rangle \\ \langle P_{\nu_\mu \rightarrow \nu_e} \rangle & \langle P_{\nu_\mu \rightarrow \nu_\mu} \rangle & \langle P_{\nu_\mu \rightarrow \nu_\tau} \rangle \\ \langle P_{\nu_\tau \rightarrow \nu_e} \rangle & \langle P_{\nu_\tau \rightarrow \nu_\mu} \rangle & \langle P_{\nu_\tau \rightarrow \nu_\tau} \rangle \end{pmatrix}. \quad (66)$$

Note that all rows and columns of M must sum to unity (total probability 1).

Using M we can describe the expected neutrino flavor content at a great distance from a neutrino source (e.g., the sun or a supernova) as follows [40]

$$\{D_e, D_\mu, D_\tau\} = M\{B_e, B_\mu, B_\tau\}, \quad (67)$$

where $\{ \}$ signifies column vectors, and D_α is the number of *detected* neutrinos of definite flavor ν_α , and B_α their number at “birth” at some distant neutrino source. Note that because neutrinos are assumed to be conserved, the total number of neutrinos at birth equals their number upon “detection,” namely,

$$B_e + B_\mu + B_\tau = D_e + D_\mu + D_\tau. \quad (68)$$

3.5.2 Proposed topological constraints on 3-flavor neutrino mixing

Given that the ν_e (ν_μ or ν_τ) neutrino flavor has the associated topology of a cylinder (Möbius strip) with respect to the internal transformation $\mathbf{F}(v)$, and assuming that topological constraints are the *primary* determinants of the matrix M describing long-distance neutrino mixtures, the matrix M is easily determined. To accomplish this we need only apply the following very general principle to neutrino-neutrino transitions:

All other things being equal, any neutrino flavor ν_α (i.e., ν_e , ν_μ or ν_τ), which undergoes neutrino-neutrino transitions that change the associated neutrino topology, will tend to be suppressed, while neutrino-neutrino transitions that maintain the associated neutrino topology will tend to be (relatively) enhanced [38].

To this principle we add the following corollary,

All other things being equal, because the ν_μ and ν_τ neutrinos have the same associated topology, they will act the same way in all neutrino-neutrino transitions (involving long-distance neutrino mixtures).

Given these principles, and assuming as stated previously that topological constraints are the *primary* determinants of the matrix M , we immediately have the following *topological* constraints on long-distance neutrino mixtures:

A. No matter what neutrino flavor (ν_α) and associated topology one starts with at some distant source (say the sun or a supernova), by the time the neutrino mixture reaches its “equilibrium” state (where all time-dependent oscillations have “damped out”), it should contain *equal* fractions of ν_μ and ν_τ , because these neutrinos have the *same* associated topology.

B. Because the ν_μ and ν_τ neutrinos have the *same* associated topology, if one starts out with *either* a pure ν_μ or a pure ν_τ source, one should end up with the *same* long-distance equilibrium mixture of ν_e , ν_μ and ν_τ .

C. If topology is the controlling factor in describing long-distance neutrino mixtures, then *there should be absolutely no (effective) difference between the two functions (of U -matrix mixing parameters), which describe $\langle P_{\nu_e \rightarrow \nu_\mu} \rangle$ and $\langle P_{\nu_e \rightarrow \nu_\tau} \rangle$, because the ν_μ and ν_τ*

neutrinos are associated with the *same* topology. Very loosely speaking we are assuming that these mathematical functions are effectively “topological invariants” with respect to the exchange of flavor indices μ and τ (see Ref. 47, pp. 20 and 21). That is, not only are the two functions $\langle P_{\nu_e \rightarrow \nu_\mu} \rangle$ and $\langle P_{\nu_e \rightarrow \nu_\tau} \rangle$ required to be *equal*, but they are also required to be equal, *term-by-term* (i.e., they are required to be *term-wise* equal). Similarly the three functions $\langle P_{\nu_\mu \rightarrow \nu_\mu} \rangle$, $\langle P_{\nu_\mu \rightarrow \nu_\tau} \rangle$ and $\langle P_{\nu_\tau \rightarrow \nu_\tau} \rangle$ are required to be term-wise equal.

Constraints A), B) and C), together with Eqs. (67) and (68), dictate that the symmetric matrix M describing long-distance neutrino mixtures must have the form expressed by

$$\begin{pmatrix} D_e \\ D_\mu \\ D_\tau \end{pmatrix} = \begin{pmatrix} a & b & b \\ b & c & c \\ b & c & c \end{pmatrix} \begin{pmatrix} B_e \\ B_\mu \\ B_\tau \end{pmatrix}. \quad (69)$$

3.5.3 Determination of the matrix M

When the proposed topological constraints of Section 3.5.2 are applied to the conventional description of neutrino mixing (62), the matrix M in (66), (67) and (69), is *uniquely* determined. To see how this happens consider the following time-average probabilities (see Appendix B in Ref. 13 for details)

$$\langle P_{\nu_e \rightarrow \nu_\mu} \rangle = 2c_1^2 s_1^2 c_2^2 + 2s_1^2 s_3^2 c_3^2 (s_2^2 - c_1^2 c_2^2) + 2s_1^2 s_2 s_3 c_1 c_2 c_3 \cos \delta (s_3^2 - c_3^2), \quad (70)$$

and

$$\langle P_{\nu_e \rightarrow \nu_\tau} \rangle = 2c_1^2 s_1^2 s_2^2 + 2s_1^2 s_3^2 c_3^2 (c_2^2 - c_1^2 s_2^2) - 2s_1^2 s_2 s_3 c_1 c_2 c_3 \cos \delta (s_3^2 - c_3^2). \quad (71)$$

According to the proposed topological constraints of Section 3.5.2, these two time-average probabilities must be *term-wise equal*, and *nonzero*. These requirements place *three* constraints on the U -matrix mixing parameters, namely, $s_1^2 > 0$, $s_2^2 = c_2^2 = \frac{1}{2}$, and

$$s_1^2 s_2 s_3 c_1 c_2 c_3 \cos \delta (s_3^2 - c_3^2) = 0. \quad (72)$$

Note that (70) and (71) are term-wise equal as required, even with the minus sign preceeding the last term of (71) because this term is of zero magnitude.

Next, the topological constraint of Section 3.5.2, namely, $\langle P_{\nu_\mu \rightarrow \nu_\mu} \rangle = \langle P_{\nu_\tau \rightarrow \nu_\tau} \rangle$ can be realized provided the U -matrix mixing parameters are further constrained by $s_3^2 = c_3^2 = \frac{1}{2}$, which also happens to satisfy (72). That is, given $s_2^2 = c_2^2$ and $s_3^2 = c_3^2$ one has (see Appendix B in Ref. 13 for details)

$$\langle P_{\nu_\mu \rightarrow \nu_\mu} \rangle = s_1^4 c_2^4 + 2c_2^4 c_3^4 (c_1^2 + 1)^2 + 8c_1^2 c_2^4 c_3^4 \cos^2 \delta, \quad (73)$$

and

$$\langle P_{\nu_\tau \rightarrow \nu_\tau} \rangle = s_1^4 c_2^4 + 2c_2^4 c_3^4 (c_1^2 + 1)^2 + 8c_1^2 c_2^4 c_3^4 \cos^2 \delta. \quad (74)$$

Now the topological constraints of Section 3.5.2 also require that $\langle P_{\nu_\mu \rightarrow \nu_\tau} \rangle$ be equal to (73) and (74). This places additional constraints on M , and the U -matrix mixing parameters. Comparing (73) and (74) with the following expression from Appendix B in Ref. 13 for $\langle P_{\nu_\mu \rightarrow \nu_\tau} \rangle$, namely,

$$\langle P_{\nu_\mu \rightarrow \nu_\tau} \rangle = s_1^4 c_2^4 + 2c_2^4 c_3^4 (c_1^2 + 1)^2 - 8c_1^2 c_2^4 c_3^4 \cos^2 \delta, \quad (75)$$

we see that the requisite equality $\langle P_{\nu_\mu \rightarrow \nu_\mu} \rangle = \langle P_{\nu_\tau \rightarrow \nu_\tau} \rangle = \langle P_{\nu_\mu \rightarrow \nu_\tau} \rangle$ leads to the constraint (note that $c_2^4 = c_3^4 > 0$)

$$c_1^2 \cos^2 \delta = 0. \quad (76)$$

Assuming [40] that $\sin^2 \delta \neq 1$ or $\cos^2 \delta \neq 0$ (i.e., CP violation is *not* maximal), we further determine from (76) that

$$c_1^2 = 0 \text{ and } s_1^2 = 1. \quad (77)$$

Note that (73), (74) and (75) exhibit the requisite term-wise equality [see item C) in Sec. 3.5.2] even with the minus sign preceeding the third term of (75), because this term is of zero magnitude.

Gathering together the predicted mixing-parameter constraints, namely, $c_2^2 = s_2^2$, $c_3^2 = s_3^2$, $s_1^2 = 1$ and $c_1^2 = 0$, we can express six of the nine time-average probabilities associated with the (symmetric) matrix M as follows (see Ref. 13).

$$\langle P_{\nu_e \rightarrow \nu_e} \rangle = 1 - 2c_3^4, \quad (78)$$

$$\langle P_{\nu_e \rightarrow \nu_\mu} \rangle = \langle P_{\nu_e \rightarrow \nu_\tau} \rangle = 2c_2^2 c_3^4, \quad (79)$$

and

$$\langle P_{\nu_\mu \rightarrow \nu_\mu} \rangle = \langle P_{\nu_\tau \rightarrow \nu_\tau} \rangle = \langle P_{\nu_\mu \rightarrow \nu_\tau} \rangle = c_2^4 (1 + 2c_3^4). \quad (80)$$

But, the previous arguments have shown that the U -matrix mixing parameters satisfy $c_2^2 = c_3^2 = \frac{1}{2}$, which leads to the following specific numerical *predictions*

$$\langle P_{\nu_e \rightarrow \nu_e} \rangle = \frac{1}{2} = a, \quad (81)$$

$$\langle P_{\nu_e \rightarrow \nu_\mu} \rangle = \langle P_{\nu_e \rightarrow \nu_\tau} \rangle = \frac{1}{4} = b, \quad (82)$$

and

$$\langle P_{\nu_\mu \rightarrow \nu_\mu} \rangle = \langle P_{\nu_\tau \rightarrow \nu_\tau} \rangle = \langle P_{\nu_\mu \rightarrow \nu_\tau} \rangle = \frac{3}{8} = c. \quad (83)$$

Given that M is symmetric, note that (81), (82) and (83) are *the only matrix elements, which could be consistent with the proposed topological constraints of Section 3.5.2, and the requirement that $\sin^2 \delta \neq 1$ or $\cos^2 \delta \neq 0$.*

Employing Eqs. (69) and (81–83), one finally has the *prediction* [40, 43]

$$M = \frac{1}{8} \begin{pmatrix} 4 & 2 & 2 \\ 2 & 3 & 3 \\ 2 & 3 & 3 \end{pmatrix}. \quad (84)$$

To summarize, *the proposed (qualitative) topological constraints on the matrix M (see Sec. 3.5.2) result in quantitative constraints on the U -matrix mixing parameters (see Eq. 62), namely, $(s_2^2 = c_2^2 = c_3^2 = s_3^2 = \frac{1}{2}, s_1^2 = 1 \text{ or } c_1^2 = 0 \text{ with } \sin^2 \delta \neq 1 \text{ or } \cos^2 \delta \neq 0 \text{ assumed})$, which result, in turn, in a unique quantitative determination of the matrix M (see Eq. 84).*

3.5.4 Topology-maintaining and topology-changing influences in equilibrium?

Not only is the matrix M in (84) a *unique* solution to the proposed topological constraints of Section 3.5.2 with $\sin^2 \delta \neq 1$ or $\cos^2 \delta \neq 0$, but it also has some very special properties that may eventually help reveal the deeper *dynamical* significance of this matrix. Note that (66), (69), and (84) have the very special property

$$\langle P_{\nu_e \rightarrow \nu_e} \rangle = \langle P_{\nu_e \rightarrow \nu_\mu} \rangle + \langle P_{\nu_e \rightarrow \nu_\tau} \rangle. \quad (85)$$

This equation says that the time-average probability that the associated ν_e topology *doesn't* change, namely,

$$P_{NC} = \langle P_{\nu_e \rightarrow \nu_e} \rangle = a, \quad (86)$$

and the time-average probability that the associated ν_e topology *does* change, namely,

$$P_C = \langle P_{\nu_e \rightarrow \nu_\mu} \rangle + \langle P_{\nu_e \rightarrow \nu_\tau} \rangle = (1 - a), \quad (87)$$

are *equal*, namely,

$$P_C = P_{NC}. \quad (88)$$

Now this equality looks very much like an “equilibrium” condition between those underlying physical influences that would act to *change* the associated ν_e topology (quantum fluctuations), and those underlying physical influences that would act to *maintain* the associated ν_e topology (e.g., topological energy “barriers”).

We will now provide further support for this proposal. In particular, consider the “joint” probability [48]

$$P = P_C \cdot P_{NC}, \quad (89)$$

and notice that

$$\frac{dP}{da} = P_C \frac{dP_{NC}}{da} + P_{NC} \frac{dP_C}{da}. \quad (90)$$

From (86) and (87) this last equation reduces to

$$\frac{dP}{da} = P_C - P_{NC}. \quad (91)$$

And, taking the second derivative, we also find

$$\frac{d^2P}{da^2} = -2 < 0. \quad (92)$$

Therefore, when $P_C = P_{NC}$, (91) and (92) tell us that the joint probability $P = P_C \cdot P_{NC}$ is a *maximum*, namely, it characterizes some *most probable* condition or “state.” And, this of course is the very essence of an “equilibrium” condition [12]. However, it must be understood that this hypothetical (long-distance) “equilibrium” between topology-changing, and topology-maintaining physical influences, is only a (cumulative) *result* of *deeper*, and largely unknown (short-distance), dynamical processes in the vacuum, which first begin to *act on neutrinos at their source—thereby, eventually establishing the equilibrium condition* $P_C = P_{NC}$ —*on time scales very much shorter than the time it takes for the time-dependent oscillations in neutrino mixtures to “damp out.”* This is an essential requirement if these hypothetical short-distance processes are to be responsible for “selecting” the (constant) U -matrix mixing parameters *prior* to neutrino mixing.

3.5.5 The current experimental situation in regard to 3-flavor neutrino mixtures

The most recent analysis [49] of Super Kamiokande and SNO data on solar neutrinos [43] is in good agreement with the prediction given by (84). If this prediction is eventually verified by further detailed observations of neutrinos from distant astronomical sources (e.g., the sun or a supernova), and/or in long-baseline terrestrial experiments (e.g., see a description of proposed Kamland experiments in Ref. 50), this will provide qualitative support for the new description of fundamental fermions, which requires, among other things, that the ν_e and (ν_μ or ν_τ) neutrino flavors start, and end, “life” as *topologically distinct* quantum objects (see Appendix A in Ref. 13, and Refs. 9, 38, 47). However, while it is conceivable that experiments could still *falsify* (84), together with the assumption [40] that CP violation is *not* maximal (i.e., $\sin^2 \delta \neq 1$ or $\cos^2 \delta \neq 0$), experimental verification of these things would *not* confirm the new description or the proposed *topological* constraints of Section 3.5.2 [51]. A much better theoretical understanding of the dynamical significance of the new 2-space description of quarks and leptons, in terms of the new internal 2-vector coordinates, will be required before one can be confident that, if finally verified, (84) really is a reflection of new physics, e.g., underlying topological constraints [51]. Clearly, additional (crucial) experimental consequences of the 2-space must be found if it is to be considered a serious candidate for a description of nature.

3.6 Does the 2-space imply the Standard Model?

Perhaps the best evidence we have for a “layer” of new physics—located somewhere between the unification scale, and the region of applicability of the standard model of particle physics—is simply that by constraining the possibilities, the 2-space description seems to *imply* something very similar to the standard model. Consider the following facts in support of this hypothesis.

First, and foremost, we have found that owing to the 2-space parameter $v = \ln M_c$, where M_c is a measure of the strong-color “multiplicity,” a symmetry associated with strong interactions such as $SU(3)_{\text{color}}$ seems to be more-or-less implicit in the 2-space description. Second, because the 2-space determines that there are some 48 fundamental fermions and antifermions [17], together with their associated global charges—identified with the global charges associated with the accidental symmetries of the strong-electroweak Lagrangian—the form of the Lagrangian is severely restricted. Third, it is clear that the 2-space dictates that we are dealing here with *flavor doublets* of quarks or leptons. And, this flavor doublet *structure* imposed by the 2-space, necessarily, severely restricts the possible *interactions* of fundamental fermions [52]. For example, we know from the 2-space description that the magnitude of the difference in electric charges between the “up” and “down” flavors within a flavor doublet is the fundamental unit of electric charge. And, if there are transitions that change an “up” flavor to a “down” flavor or vice versa, there must exist a weak intermediate boson (with unit electric charge) to accomplish this. Thinking along these lines soon leads to the notion of weak isospin, and $SU(2)_L$ flavor doublets carrying weak isospin. Fourth and finally, because the electric charge of flavor-doublet members is “explained” by the 2-space description, compatibility with electromagnetism, i.e., there is only *one* photon associated with flavor-doublet members (see Ref. 3, pp. 342–343), inevitably leads to the concept of weak hypercharge and $U(1)_Y$. So, while the 2-space certainly does *not* rigidly determine these things, it does seem to severely limit the possibilities.

In this very general way, it seems that the 2-space, and especially whatever underlying physics is responsible for it, ultimately *determines* the current form assumed by the standard model of particle physics based on $SU(3)_{\text{color}} \times SU(2)_L \times U(1)_Y$. However, until the underlying dynamics responsible for the 2-space is fully understood, the foregoing connections will remain speculative in nature.

4.0 Summary and Conclusions

It is widely believed that the explanation for fundamental-fermion (quark and lepton) flavors, flavor doublets, and family replication is to be found in theories of quantum gravity (e.g., superstrings). And, yet, as demonstrated in this paper and elsewhere [10], an analytic continuation of a Hermitian matrix used to represent the fermion-number operator, leads to a new *internal* description of quarks and leptons—a “layer” of new physics located *some-where* between the unification scale, and the region of applicability of the standard model of particle physics—that also explains such things as family replication. The new description

[9–13] involves an (internal) 2-space, and two new associated 2-vector fermion observables $\mathbf{U}$ and $\mathbf{V}$ (linearly-independent flavor-defining “coordinates”) that are somewhat analogous to the spin coordinate. And, just as the spin coordinate explains certain subtleties of atomic spectra (e.g., the fine structure) these new 2-vector coordinates explain certain subtleties of fundamental-fermion “spectra.” In particular, using the two new flavor-defining fermion coordinates $\mathbf{U}$ and $\mathbf{V}$, we have discovered, among other lesser things, that:

1) The so-called “accidental” symmetries of the Lagrangian describing strong and electroweak interactions (of fundamental fermions) are *not* accidental, but instead seem to be imposed on these Lagrangians by an underlying non-Euclidean 2-space—or whatever underlying dynamics is responsible for the 2-space, its quantization, and its associated selection rules. In particular, a central conclusion of this work is this:

The global (flavor-defining) charges associated with the “accidental symmetries” of the Lagrangian describing strong and electroweak interactions of fundamental fermions, and the global (flavor-defining) charges associated with the non-Euclidean 2-space, are (essentially) one and the same charges.

2) Family replication is apparently little more than the number of different physically acceptable ways—just *three* ways are predicted—that the “electric-charge” vector coordinate $\mathbf{Q}$, for both quark and lepton flavor doublets, can be *resolved* in the 2-space into pairs of linearly independent 2-vector coordinates $\mathbf{U}$ and $\mathbf{V}$ (i.e., $\mathbf{Q} = \mathbf{U} + \mathbf{V}$). However, while this description is reasonable, the underlying reasons for this “quantization” remain essentially unknown.

3) Unlike the situation in the standard model, particles such as the u , c and t quarks are characterized by certain associated internal *topological* differences. Similar topological differences between different neutrino flavors may help explain recent observations of (nearly) bi-maximal $\nu_\mu - \nu_\tau$ mixing. And, this in turn, *may be yet another indirect experimental indication of new dynamics (related, in part, to topology) underlying the new 2-space description of quarks and leptons.*

Finally, because the (global) accidental symmetries of the Lagrangian are at least partly determined by the 2-space, and because local symmetries such as $SU(3)_{\text{color}}$, $SU(2)_L$ and $U(1)_Y$ are also more-or-less *implicit* in the 2-space description, this description tends to imply the standard model of particle physics based on $SU(3)_{\text{color}} \times SU(2)_L \times U(1)_Y$. However, while progress has been made in making these connections, and in understanding the bewildering number of fermion flavors, two of the most important questions remain unanswered. First, what is the detailed nature of the “layer” of new physics, which is supposed to be responsible for the 2-space, its quantization, and its associated selection rules? Second, how does this “layer” of new physics arise or *emerge* from physics at the unification scale?

5.0 Acknowledgments

I wish to thank R. Zannelli for numerous helpful discussions, and for encouraging me to include footnote [25] on the possible existence (or nonexistence) of exotic fractionally-charged “quarks.” I also wish to thank M. Sheetz for skillful manuscript preparation, and for submitting the manuscript to the e-Print archive.

6.0 References and Footnotes

- [1] C. Quigg, *Gauge Theories of the Strong, Weak and Electromagnetic Interactions*, The Benjamin/Cummings Publishing Co., Reading, Mass., 1983.
- [2] K. Huang, *Quarks, Leptons and Gauge Fields*, World Scientific Publishing Co., Singapore, 1982.
- [3] O. Nachtmann, *Elementary Particle Physics—Concepts and Phenomena*, Springer-Verlag, Berlin, 1990.
- [4] M. B. Green, J. H. Schwarz and E. Witten, *Superstring Theory, Vols. 1 and 2*, Cambridge University Press, 1987.
- [5] J. M. Maldacena, “Gravity, Particle Physics and Their Unification,” arXiv:hep-ph/0002092.
- [6] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phys. Lett., B429*, **263** (1998) and arXiv:hep-ph/9803315.
- [7] S. Weinberg, *The Quantum Theory of Fields, Vol. I, Foundations*, Cambridge University Press, New York, 1995, pp. 529–531; *The Quantum Theory of Fields, Vol. II, Modern Applications*, Cambridge University Press, New York, 1996, p. 155.
- [8] S. Goudsmit and G. Ulenbeck, *Physica* **5**, 266, 1925; *Nature* **117**, 264, 1926. While flavor-defining fermion coordinates ($\mathbf{Q}$, $\mathbf{U}$, $\mathbf{V}$) are *somewhat* analogous to spin angular momentum vectors or spin “coordinates” (i.e., the total angular momentum vector $\mathbf{J}$), there is one very important difference. All spin $\frac{1}{2}$ flavors or antiflavors carry the *same* $\mathbf{J}$ and $\mathbf{J}^2 = 1/2(1/2 + 1)\hbar^2$. That is, neither $\mathbf{J}$ nor $\mathbf{J}^2$ distinguish between flavors or antiflavors whereas, as the term “flavor-defining” implies, every flavor or antiflavor is generally (but not always) associated with a *different* set of flavor-defining fermion coordinates ($\mathbf{Q}$, $\mathbf{U}$, $\mathbf{V}$). These differences, between $\mathbf{J}$ and ($\mathbf{Q}$, $\mathbf{U}$, $\mathbf{V}$), insure that while $\mathbf{J}$ vectors for *any* number of flavors and antiflavors can be *superimposed* (added), *one cannot superimpose flavor-defining fermion coordinates for two or more flavors and antiflavors* (see Sec. 3.2 in the main text for further discussion). For, if one could do this, these coordinates could not be self-consistently used to *define* individual “isolated”

flavors. In other words, flavor-defining fermion coordinates *define* (isolated) flavors or antiflavors, *not* collections of arbitrary numbers of flavors and antiflavors.

- [9] Gerald L. Fitzpatrick, *The Family Problem-New Internal Algebraic and Geometric Regularities*, Nova Scientific Press, Issaquah, Washington, 1997. Additional information: <http://physicsweb.org/TIPTOP/> or <http://www.amazon.com/exec/obidos/ISBN=0965569500>.
- [10] G. L. Fitzpatrick, “Continuation of the Fermion-Number Operator and the Puzzle of Families,” arXiv:physics/0007038 v1, 13 Jul 2000.
- [11] G. L. Fitzpatrick, “Electric Charge as a Vector Quantity,” arXiv:physics/0011073 v1, 30 Nov 2000.
- [12] G. L. Fitzpatrick, “Topological Constraints on Long-Distance Neutrino Mixtures,” arXiv:physics/0007039 v1, 13 Jul 2000. The proposal for such an equilibrium condition was first put forward in this paper, which is a relatively crude “precursor” to Ref. 13 in the present list of references.
- [13] G. L. Fitzpatrick, “Evidence for Internal Topological Constraints on Neutrino Mixtures,” arXiv:physics/0108009 v1, 7 Aug 2001.
- [14] P. A. M. Dirac, *The Principles of Quantum Mechanics* (Fourth Edition revised), Oxford University Press, 1976.
- [15] We should remind the reader how flavors are *distinguished* from one another in the standard model of particle physics. Except for mass differences, the strong and electroweak interactions of quarks are identical. Hence, from the point of view of their *interactions* alone, all quarks (leptons) are effectively *identical*. Of course, these particles are *not* identical, first because the different flavors of quarks and leptons (in principle including neutrinos) can, in fact, be *distinguished* by their masses. They can also be distinguished by certain Lorentz 4-scalar “charges,” i.e., certain global charge-conjugation-reversing (**C**-reversing) quantum numbers. These charges, being global in nature, do *not* in general couple to fields of force like color charges do. They include charges such as *baryon* and *lepton* number, *strangeness*, the *third component of global isospin*, *muon* and *electron* numbers, and other similar global *flavor-defining* charges. While it is certainly true that these flavor-defining charges can be associated with the so-called *accidental symmetries* of the Lagrangian describing strong and electroweak interactions, [e.g., $SU(3)_{\text{flavor}}$], we argue in this paper that *these charges can also be thought of as arising at a more primitive (more fundamental) level*, “below” or “beyond” the conventional standard model structure.
- [16] When weak interactions are “turned off,” flavor eigenstates and mass eigenstates are one and the same. For the most part, when we speak here of flavor eigenstates, we are referring to the situation where flavor eigenstates and mass eigenstates are the same.

- [17] Not counting spin states, we have 3 quark flavor doublets in three colors each, which yields $3 \times 2 \times 3 = 18$ states, while 3 lepton flavor doublets in one “color” each (i.e., the strong-color “multiplicity” is one), yields $3 \times 2 \times 1 = 6$ states for a total of 24 matter particles. Combining these 24 matter particles with their 24 (distinguishable) antimatter counterparts, yields a total of 48 fundamental fermions and antifermions. Not counting spin or color states, we have a total of 12 flavor eigenstates and 12 antiflavor eigenstates, for a total of 24 flavor and antiflavor eigenstates of fundamental fermions.
- [18] We assume that, because there are only 13 known force-mediating fundamental bosons (i.e., not counting spin states, we have the *photon*, the *graviton*, 8 *gluons* and 3 *weak intermediate vector bosons*—none of which carry a *flavor* index, and some of which are their *own* antiparticles) in comparison with the 48 known fundamental fermions and antifermions, the force-mediating bosons do *not* carry the new 2-vector flavor-defining coordinates. For if they did, we would naturally expect to see many more force-mediating bosons than we do. While there are, no doubt, much deeper reasons for this state of affairs, this “experimental” evidence is a reasonable indication that such flavor-defining coordinates are at least a possibility in the fermion, but *not* in the boson sector.
- [19] If there are supersymmetries, it is natural to assume that the flavor-carrying boson superpartners (squarks and sleptons) to the existing flavor-carrying fundamental fermion matter-particles (i.e., quarks and leptons), will *also* carry the new 2-vector coordinates **Q**, **U** and **V**, but the flavorless fermion superpartners (e.g., photinos, gluinos) to the existing flavorless force-mediating bosons (e.g., photons and gluons) will *not*.
- [20] J. Bernstein, *Elementary Particles and Their Currents*, W. H. Freeman and Co., San Francisco, 1968, pp. 23–25.
- [21] T. D. Lee, *Particle Physics and Introduction to Field Theory Vol. I*, Harwood Academic Publishers, New York, 1981, pp. 210–211.
- [22] Even though $\mathbf{F}(v)_{\text{diag}} = \mathbf{F}(\text{op}) = \sigma_z$, these two matrices act on entirely *different* spaces, since $\mathbf{F}(\text{op})$ is associated with the *constant* $\cos \theta = 1$, and the *variable* phase-factor $e^{-i\phi}$, whereas $\mathbf{F}(v)$ is associated with the *variable* $\cos \theta = \cosh v \geq 1$, and the *constant* phase-factor $e^{+i\phi} = +i$.
- [23] The term “non-Euclidean” is usually reserved for the geometry of *curved* spaces. However, to avoid confusing the *flat* “Lorentzian” 2-D geometry (and *discrete* transformations therein) with the *flat* Lorentzian 4-D geometry of spacetime (and *continuous* transformations therein), we prefer to use a term other than “Lorentzian” to describe the 2-space. In particular, because this 2-space is *not* Euclidean, we choose to break with tradition, and refer to this *flat* space as being “not-Euclidean,” or more correctly, *non-Euclidean*.
- [24] O. Bär and U. J. Wiese, “Can One See the Number of Colors?”, arXiv:hep-ph/0105258 v1, 24 May 2001. These authors show that in low-energy experiments involving only

pions and photons, it is impossible to “see” the number of quark colors M_c . In particular, they show that the decay width of the $\pi^0 \rightarrow \gamma\gamma$ is *not*, as is commonly claimed, proportional to M_c^2 . They further show that in such low-energy situations other values of M_c including $M_c = 5, 7, 9, \dots$, are just as reasonable as $M_c = 3$. However, when all available experimental data is properly taken into account, the number of quark colors is found to be the widely accepted number $M_c = 3$.

- [25] While it seems certain that $M_c = 1(3)$ characterizes *ordinary* leptons (quarks) it is *not* at all certain that other *extraordinary* or “exotic” values of M_c , namely, $M_c = 5, 7, 9, \dots$, are physically excluded. While we may object to such values on the grounds that they lead to as yet unseen “quarks” with unusual electric charges (e.g., $\pm 2/5$), and to corresponding strong interactions based on $SU(5)_{\text{color}}$, $SU(7)_{\text{color}}$, $SU(9)_{\text{color}}$, and so on, with their associated color-neutral composites (e.g., when $M_c = 5$, 5 “quarks” would bind “strongly” to form a color-neutral “baryon”) we have found no stronger reasons than these for excluding such states. Moreover, even if *all* of these states “lived” together in the world at the *same* time, it is still possible, in principle, that triangle anomalies could be canceled using ordinary leptons ($M_c = 1$) and an infinite “tower” of “quarks” with $M_c = 3, 5, 7, 9, \dots$, in the *same* way that triangle anomalies associated with $M_c = 1$ and $M_c = 3$ alone are canceled. It is also possible that something similar to the known three-family structure could be retained. This might be accomplished by arranging for ordinary leptons ($M_c = 1$) to be placed in a particular family along with an infinite tower of “quarks,” starting with ordinary quarks ($M_c = 3$), but also including extraordinary or exotic “quarks” with $M_c = 5, 7, 9, \dots$. Admittedly, this speculative picture leads to a rather “messy” world, but since the absence of evidence for such unusual “quark” states is *not* the same thing as evidence of their absence, we cannot simply exclude this possibility out of hand. Moreover, there is *potential* theoretical justification in string theories for such exotic fractionally-charged particles. However, it is currently unknown if the proposed exotic fractionally-charged “quark” states with $M_c = 5, 7, 9, \dots$, could be consistent with *any* existing string theory, since string theories tend to predict a relative plethora of particles having *arbitrary* fractional charges. Accordingly, until much more is known about the 2-space description, and exactly how it relates to more fundamental treatments such as string theory, it currently seems unlikely that exotic “quarks” with $M_c = 5, 7, 9, \dots$, exist in nature.

- [26] Any acceptable 2×2 matrix $\mathbf{F}(v)$ possesses just *two*, *real* linearly-independent eigenvectors, call them $\mathbf{Q}$ and $\mathbf{Q}^c$, corresponding to the two *real* eigenvalues f_m and f_a , respectively. Therefore, the matrix $\mathbf{F}(v)$ can be thought of as “producing” the conventional single-particle fermion numbers f_m and f_a via the 2-space eigenvalue equations

$$\mathbf{F}(v)\mathbf{Q} = f_m\mathbf{Q}$$

and

$$\mathbf{F}(v)\mathbf{Q}^c = f_a\mathbf{Q}^c,$$

respectively.

The 2-vector $\mathbf{Q}$ (and its scalar components—the *electric* charges of quarks and leptons) describes *matter*, while the linearly-independent 2-vector $\mathbf{Q}^c$ describes its anti-matter counterpart. The superscript c on $\mathbf{Q}^c$ is merely a label signifying antimatter. It is not an exponent or a symbol for complex conjugation. As such, it signifies only that *the 2-vectors $\mathbf{Q}$ and $\mathbf{Q}^c$ are real vectors associated with (“carried by,” “representing,” etc.) individual fundamental-fermions or antifermions, respectively*, not state vectors in some Hilbert space.

- [27] It is important to point out that the various $\mathbf{C}$ -reversing 2-scalars associated with the *internal* 2-space description are taken to be Lorentz 4-scalars in an *external* spacetime setting. That is, because q_1 and q_2 are Lorentz 4-scalars in 4-D spacetime, the components of the 2-vectors $\mathbf{U}$ and $\mathbf{V}$, and the scalar products $\mathbf{Q}^2$, $\mathbf{U}^2$, $\mathbf{V}^2$, $\mathbf{U} \bullet \mathbf{V}$ are also Lorentz 4-scalars. This connection strengthens the idea that these numbers *define* flavors in a Lorentz-invariant way.
- [28] S. Coleman and J. Mandula, “All Possible Symmetries of the S -Matrix,” Phys. Rev., Vol. 159, No. 5, 1967; S. Weinberg, *The Quantum Theory of Fields, Vol. III Supersymmetry*, Cambridge University Press, New York, 2000, pp. 2, 12–21. According to the Coleman-Mandula theorem, the possible symmetries of the S -matrix include ordinary *internal symmetries* whose generators B_k are independent of energy-momentum and spin. In other words, these group generators are Lorentz 4-“scalars.” In the present paper our internal 2-scalars—these are components, or scalar products, of the new flavor-defining fermion coordinates $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$ —are also assumed to be Lorentz 4-scalars. These 2-scalars are identified with certain flavor-defining “charges” such as *baryon number*, *lepton number*, *strangeness*, *charm* and so on. In this way, the new description honors the spirit of the Coleman-Mandula theorem, even though the internal algebraic and geometric structure we propose is, in most cases, quite different from the internal symmetries referred to in the Coleman-Mandula theorem. Our description is, if you will, a kind of “evasion” of the Coleman-Mandula theorem without violating this theorem.
- [29] The term “underlying dynamics” or “dynamics” as employed in this paper, refers to quantum dynamics supplemented by whatever “layer” of new physics (e.g., topological “energy barriers”) that may lie between the unification scale, and the region of applicability of the standard model of particle physics.
- [30] Strictly speaking, besides the specification of global flavor-defining charges, the *overall* quantum state of a fundamental fermion would, necessarily, involve a specification of the spin state, the energy-momentum state and so on, together with a specification of the particular mix of local color (gauge)-charges R , W , B , G and Y carried by each fundamental fermion. This color-mix would be determined, in turn, by something like a complementary, local $SU(5)$ color-dependent gauge description.

- [31] S. Perlis, *Theory of Matrices*, Dover Publications Inc., New York, 1991, p. 31. By definition, in any linear vector 2-space, a 2-vector such as $\mathbf{Q}$ can always be resolved into a *pair* (no more, or less) of linearly independent vectors $\mathbf{U}$ and $\mathbf{V}$ as $\mathbf{Q} = \mathbf{U} + \mathbf{V}$.
- [32] When we say that the vectors $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$ are *observables*, we mean that their associated component “charges” are mutually-commuting simultaneous observables. Hence, all of these charge-like components can be known in principle, at the *same* time, meaning that the vectors $\mathbf{Q}$, $\mathbf{U}$ and $\mathbf{V}$ can be known *simultaneously*. Thus the vector “triad” $(\mathbf{Q}, \mathbf{U}, \mathbf{V})$ is a well defined geometric object. Moreover, since $\mathbf{Q}$, $\mathbf{U}$, $\mathbf{V}$ are each Lorentz-invariant, vector triads “appear” the *same* to all observers located in arbitrary inertial coordinate systems.
- [33] Given the row vector $\mathbf{Q}_q = (q_1, q_2)$, and the column vectors $\{1, 0\}$ and $\{0, -1\}$ we see that the component quark charges associated with $\mathbf{Q}_q$ are just the non-Euclidean “projections” of $\mathbf{Q}_q$ upon the 2-space coordinate “axes” represented by the vectors $\{1, 0\}$ and $\{0, -1\}$. In particular, using the non-Euclidean scalar product , we have

$$q_1 = (q_1, q_2)\{1, 0\}$$

and

$$q_2 = (q_1, q_2)\{0, -1\}.$$

In a similar way, and for reasons of internal consistency, a hypothetical vector such as $3\mathbf{Q}_q$ would have to be projected upon the *same* axes $\{1, 0\}$ and $\{0, -1\}$ to obtain the component charges. However, only the charge projections $3q_1$ and $3q_2$ are obtained in this case. Other charges such as $q = (2q_1 + q_2)$ *cannot be obtained by means of such projections*.

- [34] Given the 2 by 2 matrix $\mathbf{R}$ in Eq. (51) in the main text, we see immediately that the effect of $\mathbf{R}$ on the column vectors $\{1/2, -1/2\}$, $\{1, 0\}$ and $\{0, -1\}$ is simply $\mathbf{R}\{1/2, -1/2\} = \{1, 0\}$ and $\mathbf{R}\{1, 0\} = \{0, -1\}$, or $\mathbf{R}^2\{1/2, -1/2\} = \{0, -1\}$.
- [35] It is shown elsewhere (see Ref. 9, pp. 63–65 in the current reference list) that the matrices $\mathbf{R}$ and $\mathbf{R}^{-1}$ are charge-conjugation-reversing (**C**-reversing). In particular, the *components* of $\mathbf{R}$ and $\mathbf{R}^{-1}$ transform like **C**-reversing, Lorentz-invariant (4-scalar) *charges*. And, this in turn means that $\mathbf{R}$ and $\mathbf{R}^{-1}$ are *Lorentz-invariant*.
- [36] J. Rosen, *A Symmetry Primer for Scientists*, John Wiley and Sons, New York, 1983, p. 16. The symmetry-evolution principle, which is closely associated with the second law of thermodynamics, states: *For an isolated physical system the degree of symmetry cannot decrease as the system evolves, but either remains constant or increases*.
- [37] Note that the change described in $\Delta\mathbf{U} = \mathbf{U}_2 - \mathbf{U}_1$ involves the “addition” of the vectors $\mathbf{U}_2$ and $-\mathbf{U}_1$. However, this addition does *not* take place at the same spacetime

coordinate. That is, $\Delta\mathbf{U} = \mathbf{U}_2 - \mathbf{U}_1$ describes a *transition* in which the initial $\mathbf{U}$ -vector describing a particle is $\mathbf{U}_1$, and the final $\mathbf{U}$ -vector describing the particle (possibly a different particle) is $\mathbf{U}_2$. That is, this transition involves only *one* current. So, the “addition” expressed by the change $\Delta\mathbf{U} = \mathbf{U}_2 - \mathbf{U}_1$ does *not* violate our rule that like 2-vectors (i.e., $\mathbf{Q}$, $\mathbf{U}$ or $\mathbf{V}$) for two or more fundamental fermions *cannot* be added or superimposed. The same arguments obviously apply to changes described by $\Delta\mathbf{Q}$ and $\Delta\mathbf{V}$.

- [38] D. J. Thouless, *Topological Quantum Numbers in Nonrelativistic Physics*, World Scientific, Singapore, 1998. Numerous examples from physics and mathematics could be cited to establish the following very general principle:

In physical systems characterized by a well defined topology, topology-change tends to be suppressed, relative to the condition of topology-maintenance.

The basis for this principle is that topology change involves either an energy “expense” (an energy “barrier” must be overcome) or a violation of some “topological charge” conservation law, or both. In general, we expect this principle to apply to any discontinuous operation such as the *abstract* equivalent of “tearing,” and subsequently “glueing,” surfaces back together to form states with new topologies. For example, one cannot continuously deform a doughnut into a sphere. It must first be “cut” and then “glued” back together in a new way to achieve such a transformation.

Now, given the “facts” regarding the associated topology of neutrinos with respect to the (abstract) internal transformation $\mathbf{F}(v)$, and *assuming that there are no “energy barriers” associated with transitions between neutrinos having the same associated topology*, the foregoing principle suggests that these facts could have the following dynamical significance:

Associated topology change in neutrino-neutrino transitions ($\nu_\alpha \rightarrow \nu_\beta$) is suppressed by topological energy and/or topological charge “barriers,” while associated topology maintenance in neutrino-neutrino transitions is relatively enhanced.

It should be emphasized that even though the ν_μ and ν_τ neutrinos have the *same* associated topology with respect to $\mathbf{F}(v)$, they are quite *distinct* in other respects. For example, in conventional weak decays or weak capture reactions involving these particles, mu- and tau- numbers are *separately* conserved.

- [39] The baryon (lepton) asymmetry of the universe insures that B (L) are not absolutely conserved. However, violation of these conservation laws is very strongly suppressed as evidenced by the long life of the proton.
- [40] H. Georgi and S. L. Glashow, “Neutrinos on Earth and in the Heavens,” arXiv:hep-ph/9808293, and hep-ph/9808293 v2, page 5, Eq. (20). It is interesting, and probably significant that Georgi and Glashow independently arrived at Eq. (84) in the main text of the present paper (see Eq. 20 in their paper) starting from six experimental neutrino

“facts,” some of which are completely different than the facts and assumptions employed in the present paper. In particular, these authors began by assuming

1. There are just three chiral neutrino states having Majorana masses.
2. Atmospheric neutrinos rarely oscillate into electron neutrinos.
3. Atmospheric muon neutrinos suffer maximal, or nearly maximal, two flavor oscillations into tau neutrinos.
4. & 5. Two experimentally (and theoretically) motivated assumptions regarding the order of magnitude of individual neutrino masses, and various mass squared differences.

and

6. Neutrinoless double beta decay either does *not* occur or is highly suppressed.

Given these six “facts” these authors derived both the neutrino mass matrix, and partially determined the associated CKM-like unitary matrix U , which describes neutrino mixtures. From U they then derived Eq. (84) in the present paper. In closing, these authors also noted that CP-violation in this situation could be “superweak.” Clearly, this would be consistent with the assumption in the present paper that CP-violation is *not* maximal, i.e., $\sin^2 \delta \neq 1$ or $\cos^2 \delta \neq 0$.

While the present approach, and that of Georgi and Glashow in deriving Eq. (20) in their paper, are quite different they are, nevertheless, expected to be compatible. In particular, we anticipate that such things as *neutrino masses and mixing parameters, ultimately owe their existence to physics at a deeper level, where topological considerations, of the kind proposed in the present paper, are expected to play an important role.*

- [41] A. P. Balachandran, “Bringing Up a Quantum Baby,” arXiv:quant-ph/9702055.
- [42] G. Holzwarth, “Formation of Extended Topological-Defects During Symmetry-Breaking Phase Transitions in $O(2)$ and $O(3)$ Models”, arXiv:hep-ph/9901296. Analogous examples of fluctuation-induced topology-change in macroscopic objects (e.g., destruction of topological “defects” due to thermal fluctuations at phase transitions) abounds. For example, otherwise persistent (“conserved”) *topological defects* in crystals can be destroyed by raising the temperature sufficiently (i.e., by melting the crystal). Similarly, otherwise persistent (“conserved”) magnetic *flux-tubes* in Type II superconductors and/or *vortices* in a superfluid, can both be destroyed by raising the temperature above the critical temperature T_c . And, conversely, topological defects are *always* created when such macroscopic systems first condense (or crystallize) as the temperature is lowered. We imagine that something roughly similar can happen when quantum-fluctuations (vacuum fluctuations) act on otherwise very similar quantum objects (i.e., quantum objects such as neutrinos having the same electric charge, spin, lepton number, and nearly

identical mass) that also happen to start “life” as topologically-distinct quantum objects. That is, we are assuming that, if not prevented by some absolute conservation law, transitions between such states (e.g., ν_e and ν_τ neutrinos) will be *catalyzed* by quantum fluctuations.

- [43] The Super-Kamiokande, Kamiokande Collaboration, arXiv:hep-ex/9810001, and the SNO Collaboration, arXiv:nucl-ex/0106015.
- [44] M. Gronau, “Patterns of Fermion Masses, Mixing Angles and CP Violation,” in *The Fourth Family of Quarks and Leptons, First International Symposium*, edited by D. B. Cline and Amarjit Soni, Annals of The New York Academy of Sciences, New York, Volume 518, 1987, p. 190. According to the scheme described in the present paper, the fundamental fermions $d, \nu_e(s, b, \nu_\mu, \nu_\tau)$ have the topology associated with a cylinder (Möbius strip) with respect to the internal transformation $\mathbf{F}(v)$. Hence, we imagine that if, in some imaginary world, quark couplings to hypothetical Higgs mass-producing fields were nearly “switched off,” the resulting “low” mass d, s and b quark mixtures (located *inside* strongly-bound quark composites) could conceivably look very much like unbound or “free” ν_e, ν_μ and ν_τ low mass neutrino mixtures—owing to topological influences of exactly the kind proposed in the present paper. In particular, in this imaginary world, we would expect the s and b quarks (like the ν_μ and ν_τ neutrinos, respectively) to exhibit (nearly) bi-maximal mixing. However, because of uncertainty-principle considerations, and because the b quark mass is, in the “real” world, very much *larger* than the d and s quark masses, very little mixing with the b quark occurs in d, s, b quark mixtures *in spite of topological influences that would tend to encourage this*. Thus, in the case of strongly-interacting quarks, it seems very likely that topological constraints of the kind considered in the present paper can, at most, play a *minor* role in determining such things as CKM-type matrix elements. For example, in the case of d and s quarks, Gronau shows that the mixing angle θ_c depends on quark masses since the matrix element V_{12} , and the d, s quark masses m_d and m_s , respectively, are known to be empirically related via $V_{12} = \sin \theta_c = \sqrt{\frac{m_d}{m_s}}$, where $m_s > m_d \neq 0$. This is nothing like the corresponding matrix element for ν_e and ν_μ neutrinos found in the present paper. Thus, although a detailed understanding of these matters is far from being achieved, it appears that topological influences are allowed to play a major role in neutrino, but *not* in quark mixing, because of the small value, and near degeneracy of neutrino masses in comparison to the large differences between d, s and b quark masses.
- [45] Although conventional treatments of neutrino mixing do not involve topological considerations, they do describe mixing among neutrinos that initially have a *definite flavor*. And, because we argue that a neutrino with a definite flavor is also associated with a *definite topology*, hereinafter we will often use these terms together as in “... neutrinos have a definite flavor and associated topology.”

- [46] Ta-Pei Cheng and Ling-Fong Li, *Gauge Theory of Elementary Particle Physics*, Clarendon Press, Oxford, 1984, pp. 409–414. While this reference provides a good summary of the formalism involved in neutrino mixing, the reader should be cautioned that there are several “typographical” errors that could cause confusion. For example, in Eq. (13.22) the first term in the expression for the matrix element U_{32} should read $c_1 s_2 c_3$ *not* $c_1 s_2 s_3$. In Eq. (13.31) the last term in the expression for the time-average probability $\langle P_{\nu_e \rightarrow \nu_\tau} \rangle$ should be preceded by a *minus* sign, *not* a plus sign. Finally, the left hand side of Eq. (13.33) should read $P_{\nu_\alpha \rightarrow \nu_\beta}$ *not* $\langle P_{\nu_\alpha \rightarrow \nu_\beta} \rangle$.
- [47] C. Nash and S. Sen, *Topology and Geometry for Physicists*, Academic Press, New York, 1983.
- [48] Note that for all practical purposes, time-dependent *quantum* probabilities $P_{\nu_\alpha \rightarrow \nu_\beta}$ become time-average or *classical* probabilities $\langle P_{\nu_\alpha \rightarrow \nu_\beta} \rangle$ when t is sufficiently large. For this reason, it makes sense to consider classical probability measures such as the “joint” probability $P = P_C P_{NC}$, where P_C and P_{NC} are time-averages of quantum probabilities.
- [49] A. Bandypadhyay et. al., “Three Generation Neutrino Oscillation Parameters after SNO,” arXiv:hep-ph/0110307 v2, 19 Dec 2001. Equation 19 in this paper gives a three-generation mixing matrix U , which leads to a matrix describing long distance equilibrium neutrino mixtures given by $M = (M_{11} = 0.6, M_{12} = M_{13} = M_{21} = M_{31} = 0.2, M_{22} = M_{23} = M_{32} = M_{33} = 0.4)$; P. F. Harrison et. al., “Tri-Bimaximal Mixing and the Neutrino Oscillation Data,” arXiv:hep-ph/0202074, 7 Feb 2002. Equation 6 in this paper gives a three-generation mixing matrix U , which leads to a matrix describing long distance equilibrium neutrino mixtures given by $M = (M_{11} = 0.56, M_{12} = M_{13} = M_{21} = M_{31} = 0.22, M_{22} = M_{23} = M_{32} = M_{33} = 0.39)$. Both of these experimentally determined matrices compare favorably with the theoretical predictions of the present paper (see Eq. 84 in the main text), namely, $M = (M_{11} = 0.5, M_{12} = M_{13} = M_{21} = M_{31} = 0.25, M_{22} = M_{23} = M_{32} = M_{33} = 0.375)$.
- [50] A. Suzuki, Eighth International Workshop on Neutrino Telescopes, Venice, 1999.
- [51] Even if the matrix M in Eq. (84) in the main text is verified by experiment, we could not be certain that the constraints expressed by items A), B) and C) in Section 3.5.2, are *topological* constraints related to the 2-space description of quarks and leptons. There are three basic reasons for this assertion.

First, we do not know why the topology of vector triads [with respect to the internal transformation $\mathbf{F}(v)$] should be *relevant* to neutrino mixing. Second, we do not have any detailed understanding of the hypothetical *mechanism* that supplies the “energy barriers” or “topological charge conservation laws” that serve to inhibit topology change in neutrino mixing. Third, as indicated in Ref. 13, Appendix A (Sec. A.4) in this list, we cannot be absolutely certain that the ν_μ neutrino is associated with the requisite second-family Möbius topology with respect to $\mathbf{F}(v)$. What we do know with certainty is that, regardless of their physical origins, the constraints of Section 3.5.2, together

with the additional assumption that $\sin^2 \delta \neq 1$ or $\cos^2 \delta \neq 0$, definitely determine M *uniquely*.

- [52] However, all of the flavor-defining charges associated with these 2-space flavor doublets are *global* in nature, so this tells us nothing directly about the *interactions* of flavor doublet members. Nevertheless, the “up”–“down” flavor doublet *structure* imposed by the 2-space, necessarily, severely restricts the possible interactions.